

NOBANIS – Invasive Alien Species Fact Sheet

Azolla filiculoides

Author of this species fact sheet

Dipl.-Biol. Andreas Hussner, Heinrich-Heine-Universität Düsseldorf, Abt. Geobotanik
Universitätsstraße 1, 40225 Düsseldorf, Tel.: 0049-211-8113683, Andreas.Hussner@uni-duesseldorf.de

Bibliographical reference – how to cite this fact sheet:

Hussner, A. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Azolla filiculoides*. – From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org, Date of access x/x/200x.

Species description

Scientific names : *Azolla filiculoides* Lam. 1783, Azollaceae

Synonyms: *Azolla caroliniana* Willd., *Azolla rubra* R. Br., *Azolla japonica* Franch. and Sav., *Azolla arbuscula* Desv., *Azolla filiculoides* var. *rubra* Lam. (R.Br.) Strasb, *Azolla magellanica* Wild., *Azolla squamosa* Molina.

Some scientists consider *A. caroliniana*, *A. japonica* and *A. rubra* as independent species.

Common names: large mosquito fern (GB), red water fern (GB), Großer Algenfarn (DE), Andemadsbregne (DK), *Azolla karolinska* (PL), šakotoji azolė (LT), Mossbräken (SE), Limaskasaniainen (FI).



Fig. 1 and 2. Single plants of *Azolla filiculoides*, photos by A. Hussner.



Fig. 3 and 4. Thin and dense mats of *Azolla filiculoides*, photos by A. Hussner.




Fig. 5. *Azolla filiculoides* plants in a heated pond in winter; the plants become red with an increasing distance from the heater, photo by A. Hussner.

Species identification

A. filiculoides is a heterosporous, up to 2.5 (10) cm large floating fern. *Azolla* plants are polygonal or triangular in shape (Lumpkin and Plucknett 1980). The sporophytes consist of two-lobed leaves and rhizomes. The lower lobes of the leaves are usually larger than the upper. Svenson (1944) described the lower lobes as so adapted for floating the plant, that only the lower surface of these lower lobes comes in contact with water.

The plants are dark green to reddish and float on the water surface, either individually or in mats, which can reach a thickness of up to 20 cm (McConnachie *et al.* 2004). A characteristic of the genus *Azolla* is the symbiotic relationship with the nitrogen-fixing blue-green alga *Anabaena azollae*. When *A. filiculoides* plants are exposed to strong sunlight they obtain a red colour. The same occurs in wintertime. In shade they always remain green (Janes 1998a, Moore 1953). Janes (1998a) described three different phenotypes of *A. filiculoides* under different habitat and climate conditions.

Native range

A. filiculoides is native to warm temperate and subtropical America through Western North America (including Alaska). West (1953) described *A. filiculoides* furthermore as a species, which in former times was native to Europe (Ekman 1998, 1999, O'Brien and Jones 2003), but died out during the last Ice Ages.

Alien distribution

History of introduction and geographical spread

A. filiculoides is actually a cosmopolitan plant, with occurrences in South Africa (Hill and Cilliers 1999, McConnachie *et al.* 2003, 2004, Gratwicke and Marshall 2001), Asia (Ahmad 1941, 1943, Kitoh *et al.* 1993), Australia, South-, Central- and North America (Svenson 1944), South-, West-, Central- and North Europe (Bernhardt 1991, Birkenbeil 1974, Ferreira *et al.* 1998, Janes 1998a, b, Kohler 1995, Hussner and Lösch 2005, West 1953) and Scandinavia (Rune and Jorgensen 1997). The species was introduced to Europe in 1880 near Bordeaux (West 1953). First plants were reported from France and since then, the species spread to nearly whole Europe with a main occurrence in Atlantic-Mediterranean regions.

Pathways of introduction

A. filiculoides is able to reach new regions by waterfowls, in ballast tanks of ships or by human influence (intentionally or unintentionally). Discharge by aquarium keepers may play an important role in spread of this species. Rune and Jorgensen (1997) described two examples of intentional releases of *A. filiculoides* plants in Denmark.

Alien status in region

A. filiculoides is an alien plant to Europe and Scandinavia (see table 1). In Germany *A. filiculoides* is common (Bernhardt 1991, Birkenbeil 1974, Kohler 1995, Jaeger and Werner 2002, Kowarik 2003, Hussner and Lösch 2005), but there are no reports on such thick mats as they are reported from other countries (McConnachie *et al.* 2003). Rune and Jorgensen (1997) described *A. filiculoides* from 11 different sites in Denmark. *A. filiculoides* is found in a few localities in Southern Sweden. It is uncertain if *A. filiculoides* is established or if occurrences are incidental. In Poland the species is known from some ephemeral locations in the SW and SE regions (Rostanski and Sowa 1986-1987, Wolkowyski 1999). In Lithuania *A. filiculoides* was recorded abundantly growing in an exploited peatbog in 1946, later became extinct (Gudžinskas 2000).

Country	Not found	Not established	Rare	local	Common	Very common	Not known
Denmark			X				
Estonia	X						
European part of Russia	X						
Finland	X						
Faroe Islands	X						
Germany					X		
Greenland	X						
Iceland	X						
Latvia	X						
Lithuania		X					
Norway							
Poland		X					
Sweden			X				

Table 1. The frequency and establishment of *Azolla filiculoides*, please refer also to the information provided for this species at www.nobanis.org/search.asp. Legend for this table: **Not found** –The species is not found in the country; **Not established** - The species has not formed self-reproducing populations (but is found as a casual or incidental species); **Rare** - Few sites where it is found in the country; **Local** - Locally abundant, many individuals in some areas of the country; **Common** - Many sites in the country; **Very common** - Many sites and many individuals; **Not known** – No information was available.

Ecology

Habitat description

A. filiculoides settles in ponds, ditches, water reservoirs, wetlands, channels and slow flowing rivers. *A. filiculoides* can be found in sunny to shady parts of the water bodies. Through its symbiotic association with *Anabaena azollae*, the floating fern is able to grow in nitrogen-deficient waters. Kitoh *et al.* (1993) observed that less phosphorus can limit the growth of the species. The species grows at best at 15-20°C and high irradiance (Janes 1998a, Tung and Watanabe 1983, Watanabe and Berja 1983). Wong *et al.* (1987) stated, that optimum temperature for both nitrogen fixation and oxygen evolution is 25°C. Janes (1998a) reported, that plants die under laboratory conditions at temperatures below – 4°C. Janes (1998a) described *A. filiculoides* as the most frost-tolerant of the *Azolla* species. Wong *et al.* (1987) reported that *A. filiculoides* withstands field temperatures of – 10 to – 15°C. Janes (1998a) observed successfully overwintered vegetative plants in Worcester (UK), where the minimum air temperature reached – 10°C. Janes (1998a) described, that plants are able to survive encasement in ice for at least one week, but these parts of the plants, which protruded above the ice, were killed.

Reproduction and life cycle

A. filiculoides has a surface-area doubling time of 7-10 days under favourable conditions. Kitoh *et al.* (1993) observed a doubling of the biomass every 2.2 – 3.4 days under laboratory conditions. The growth rate of *A. filiculoides* increased with increased photoperiod. Janes (1998a, b) investigated the sporulation and germination of *A. filiculoides* and showed that the species sporulate regularly at many sites in Britain, most often between May and November. From Germany sporulating plants are known, too (pers. obs.). It seems that sporulation of this species is regulated by the interacting effects of light intensity, photoperiod, temperature, pH and nutrient availability. Janes (1998b)

suggested that a thick mat of 8 kg m⁻² fresh biomass may produce 85000 megasporocarps and 380000 microsporocarps. In laboratory experiments a maximum of germination was reached at 20°C but at a constant temperature of 5°C no germination was observed (Janes 1998b). For germination a temperature above 10°C and light are necessary. Sub-zero temperatures of about – 10°C for at least 18 days had no influence on further germination of the sporocarps. The heterosporous life cycle of the genus *Azolla* is illustrated in Fig 6.




Fig. 6. Heterosporous life cycle of *Azolla* (redrawn from Lumpkin and Plucknett 1980)

Dispersal and spread

A. filiculoides can spread locally by waterfowls, water sport tackle (diving, fishing, water skiing etc.) or by discharge from aquarium keepers.

Impact

Affected habitats and indigenous organisms

A. filiculoides settles in ponds, ditches, water reservoirs, channels and slow flowing rivers. In some cases the species can be found together with *Lemna minuta*, *Lemna minor* and *Spirodela polyrhiza*. In other cases, *A. filiculoides* forms dense monospecific mats. These mats of floating plants can affect the water by eliminating submerged plants and algae (Janes *et al.* 1996), preventing their photosynthesis and blocking oxygen diffusion. Also populations of animals are reduced in the water beneath the mats (Gratwicke and Marshall 2001).

Genetic effects

Janes (1998b) reported, that there is some evidence that *A. filiculoides* might have adapted to the British climate since its introduction. Similar adaptations are possible for the population in other countries.

Human health effects

There are no human health effects known.

Economic and societal effects (positive/negative)

The genus *Azolla* is useful in rice fields, because it can assimilate atmospheric nitrogen gas owing to the nitrogen fixation by cyanobacteria (blue green alga) living in the cavities located at the lower side of upper (dorsal) lobes of leaf. By this, *Azolla* is used as a green manure in rice fields in Asia

since several decennia (Choudhury and Kennedy 2004, Kimura 2005, De Macale and Vlek 2004, Mandal *et al.* 1999, Nayak *et al.* 2004, Moore 1969). Aside from this fertilizing effect in cultivation of rice, *Azolla* may influence the pH of the water (Vlek *et al.* 2002) Shiomi and Kitoh (2001), Parthasarathy *et al.* (2002), Reyes and Fermin (2003), described the use of *Azolla* as a feed for animals, particularly for fishes.

In the last years the genus *Azolla* has become more and more important in wastewater treatments to eradicate different metals or to remove nitrogenous compounds from the water (Allison *et al.* 2000, Cohen-Shoel *et al.* 2002, Costa *et al.* 1999, Forni *et al.* 2001a, b, Forni *et al.* 2002, Gardea *et al.* 2005, Ghobrial and Siam 1998, Oren Benaroya *et al.* 2004, Sanyahumi *et al.* 1998, Shiny *et al.* 2004, Stepniewska *et al.* 2005, Zhao and Duncan 1998, Zhao *et al.* 1999 a, b).

On the other hand, *Azolla* can form such dense and thick mats, that it may be impossible to row a boat through it (Moore 1969). *Azolla* mats can reduce the population of different animals from water beneath the mats (Gratwicke and Marshall 2001), and furthermore submerged plants can be eradicated by shading (Janes *et al.* 1996).

Management approaches

Prevention methods

Prevention methods are not known.

Eradication, control and monitoring efforts

Control options for *A. filiculoides* are limited. Due to a surface-area doubling time of 7-10 days, mechanical control is impractical. Hill and Cilliers (1999) considered manual, mechanical and herbicidal control as undesirable and only small infestations of *Azolla* can be removed manually using fine meshed nets. Barreto *et al.* (2000) described the biological control of *Azolla* with fungi. Hill (1998), Hill and Cilliers (1999) and McConnachie *et al.* (2003, 2004) described a successful biological control of bigger occurrences by the frond-feeding weevil *Stenopelmus rufinasus* in South Africa, where *A. filiculoides* covered dams and water reservoirs.

Education and awareness

No

Knowledge and research

There is a broad knowledge about the biology and physiology of *A. filiculoides*. Since several decennia scientists around the world have investigated the genus *Azolla*. Moore (1969) and Lumpkin and Plucknett (1980) have made excellent reviews of literature. In the last years the control of *A. filiculoides* has gained more and more importance (McConnachie *et al.* 2003, 2004). In addition the research about bioaccumulation and phytoremediation has received high interest in science of wastewater treatment (Cohen-Shoel *et al.* 2002, Forni *et al.* 2001, Forni *et al.* 2002, Gardea *et al.* 2005, Ghobrial and Siam 1998, Oren Benaroya *et al.* 2004, Sanyahumi *et al.* 1998, Shiny *et al.* 2004, Stepniewska *et al.* 2005, Zhao and Duncan 1998, Zhao *et al.* 1999 a, b).

Recommendations or comments from experts and local communities

A comprehensive investigation of actual distribution and involved water bodies will be helpful for an estimation of the further spread of this species in our region.

References and other resources

Contact persons

Andreas Hussner (DE) Heinrich-Heine-Universität Düsseldorf, Department of Geobotany, Universitätsstraße 1, 40225 Düsseldorf, Tel.: 0049-211-8113683, E-mail: Andreas.Hussner@uni-duesseldorf.de

Daniel Larsson (SE) Department of Environmental Analysis, Swedish University of Agricultural Sciences, P.O. Box 7050, SE-757 55 Uppsala. Tel. +46 18 67 31 42. E-mail: daniel.larsson@ma.slu.se

Christina Birnbaum (EE) Estonian University of Life Sciences, Institute of Agriculture and Environmental Sciences, Kreutzwaldi 64 Tartu 51014, phone, E-mail: chbirnbaum@yahoo.com

Dr Dan Wolkowycski (PL) Białystok Technical University, Institute of Environmental Engineering and Protection, Division of Landscape Ecology, ul. Wiejska 45E, 15-351 Białystok, Poland, Tel. 0048 85 74 28 401 ext. 307, E-mail: danwolko@pb.bialystok.pl

Henrik Jørgensen (DK) Danish Forest and Nature Agency, Ministry of the Environment, Haraldsgade 53, DK-2100 Copenhagen Ø; Phone: +45 3947 2000, E-mail: hjg@sns.dk

Links

[Aquatic neophytes](#) – in German

Flora of Northern Ireland – [fact sheet on *Azolla filiculoides*](#)

Canadas Plant Species – [Azolla filiculoides](#) incl. distribution map

Biological Control of Red Water Fern in South Africa - [report](#)

[Azolla in USA](#)

References

Ahmad, G. (1941): Effect of light intensity and temperature on the growth of *Azolla filiculoides*. Journal of the Indian Botanical Society 20: 213-226

Ahmad, G. (1943): Interrelationship between the compensation point, temperature coefficient and growth of *Azolla filiculoides*. Journal of the Indian Botanical Society 22-23: 101-104

Allison, G., Stagnitti, F., Colville, S., Hill, J. and Coates, M. (2000): Growth of floating aquatic macrophytes in alkaline industrial wastewaters. Journal of Environmental Engineering 126: 1103-1107

Barreto, R., Charudattan, R., Pomella, A. and Hanada, R. (2000): Biological control of neotropical aquatic weeds with fungi. Crop Protection 19: 697-703

Bernhardt, K.G. (1991): Zur aktuellen Verbreitung von *Azolla filiculoides* Lam. (1783) and *Azolla caroliniana* Willd. (1810) in Nordwestdeutschland. Floristische Rundbriefe 25: 14-19

Birkenbeil, H. (1974): *Azolla filiculoides* Lamk. im Nordheimer Altrhein. Hessische Floristische Briefe 23: 14-16

- Cohen-Shoel, N., Barkay, Z., Ilzyer, D., Gilath, I. and Tel-Or, E. (2002): Biofiltration of toxic elementary by *Azolla* biomass. *Water, Air, and Soil Pollution* 135 (1-4): 93-104
- Costa, M.L., Santos, M.C. and Carrapico, F. (1999): Biomass characterization of *Azolla filiculoides* grown in natural ecosystems and wastewater. *Hydrobiologia* 415: 323-327
- De Macale, M.A.R., Vlek, P.L.G. (2004) : The role of *Azolla* cover in improving the nitrogen use efficiency of lowland rice. *Plant and Soil* 263 (1-2): 311-321
- Ekman, S.R. (1998): Pleistocene pollen stratigraphy from borehole 81/34, Devil's hole area, central North Sea. *Quaternary Science Reviews* 17: 855-869
- Ekman, S.R. (1999): Early Pleistocene pollen biostratigraphy in the central North Sea. *Review of Palaeobotany and Palynology* 105: 171-182
- Ferreira, M.T., Catarino, L. and Moreira, I. (1998): Aquatic weed assemblages in an Iberian drainage channel system and related environmental factors. *Weed research* 38(4): 291-300
- Forni, C., Chen, J., Pancioni, L. and Grilli Caiola, M. (2001a): Evaluation of the fern *Azolla* for growth, nitrogen and phosphorous removal from wastewater. *Water research* 35 (6): 1592-1598
- Forni, C., Nicolai, M.A. and D'Egidio, M.G. (2001b): Potential of the small aquatic plants *Azolla* and *Lemna* for nitrogeous compounds removal from wastewater. *Progress in Water Resources*: 315-324
- Forni, C., Cascone, A., Fiori, M. and Migliore, L. (2002): Sulphadimethoxine and *Azolla filiculoides* Lam.: A model for drug remediation. *Water research* 36 (13): 3398-3403.
- Gardea-Torresdey, J.L., Peralta-Videa, J.R., De La Rosa, G. and Parson, J.G. (2005): Phytoremediation of heavy metals and study of the metal coordination by X-ray absorption spectroscopy. *Coordination Chemistry Reviews* 249: 1797-1810
- Ghobrial, M.G. and Siam, E.E. (1998): The use of the water velvet *Azolla filiculoides* in wastewater treatment. *Journal of the Chartered Institution of Water and Environmental Management* 12 (4): 250-253
- Gratwicke, B. and Marshall, B.E. (2001): The impact of *Azolla filiculoides* Lam. on animal biodiversity in streams in Zimbabwe. *African Journal of Ecology* 39 (2): 216-218
- Gudžinskas, Z. (2000): Conspectus of alien plant species of Lithuania. 15. *Azollaceae*, *Pinaceae*, and *Salicaceae*. *Botanica Lithuanica*, **6(3)**: 235-242.
- Hill, M.P. (1998): Life history and laboratory host range of *Stenopelmus rufinusus*, a natural enemy for *Azolla filiculoides* in South-Africa. *BioControl* 43 (2): 215-224
- Hill, M.P. and Cilliers, C.J. (1999): *Azolla filiculoides* Lamarck (Pteridophyta: Azollaceae), its status in South Africa and control. *Hydrobiologia* 415: 203-206
- Hussner, A. and Lösch, R. (2005): Alien aquatic plants in a thermally abnormal river and their assembly to neophyte-dominated macrophyte stands (River Erft, Northrhine-Westphalia). *Limnologia* 35: 18-30
- Jaeger, E. and Werner, K. (2002): Rothmaler, Exkursionsflora von Deutschland. Band 4, Gefäßpflanzen, kritischer Band, 9. Auflage, Spektrum, Heidelberg/Berlin
- Janes, R. (1998a): Growth and survival of *Azolla filiculoides* in Britain. 1. Vegetative reproduction. *New Phytologist* 138: 367-376
- Janes, R. (1998b): Growth and survival of *Azolla filiculoides* in Britain. 2. Sexual reproduction. *New Phytologist* 138: 377-384

- Janes, R., Eaton, J.W. and Hardwick, K. (1996): The effects of floating mats of *Azolla filiculoides* Lam. and *Lemna minuta* Kunth on the growth of submerged macrophytes. *Hydrobiologia* 340: 23-26
- Kimura, M. (2005): Populations, community composition and biomass of aquatic organism in the floodwater of rice fields and effects of field management. *Soil Science and Plant Nutrition* 51 (2): 159-181
- Kitoh, S., Shiomi, N. and Uheda, E. (1993): The growth and nitrogen fixation of *Azolla filiculoides* Lam. in polluted water. *Aquatic botany* 46: 129-139
- Kohler, A. (1995): Neophyten in Fließgewässern – Beispiele aus Süddeutschland und dem Elsaß. *Schriftenreihe für Vegetationskunde* 27: 405-412
- Kowarik, I. (2003): *Biologische Invasionen: Neophyten und Neozoen in Mitteleuropa*. Eugen Ulmer Verlag, Stuttgart, 380 S.
- Lumpkin, T.A. and Plucknett, D.L. (1980): *Azolla: Botany, Physiology, and Use as a Green Manure*. *Economic Botany* 34 (2): 111-153
- Mandal, B., Vlek, P.L.G. and Mandal, L.N. (1999): Beneficial effects of blue-green algae and *Azolla*, excluding supplying nitrogen, on wetland rice fields: A review. *Biology and Fertility of Soils* 28 (4): 329-342
- McConnachie, A.J., Wit, M.P. de, Hill, M.P. and Byrne, M.J. (2003): Economic evaluation of the successful biological control of *Azolla filiculoides* in South Africa. *Biological Control* 28: 25-32
- McConnachie, A.J., Hill, M.P., Byrne, M.J. (2004): Field assessment of a frond-feeding weevil, a successful biological control agent of red waterfern, *Azolla filiculoides*, in southern Africa. *Biological control* 29: 326-331
- Moore, A.W. (1969): *Azolla: biology and agronomic significance*. *Botanical Review* 35: 17-35
- Nayak, S., Prasanna, R., Pabby, A., Dominic, T.K. and Singh, P.K. (2004): Effect of urea, blue green algae and *Azolla* on nitrogen fixation and chlorophyll accumulation in soil under rice.
- O'Brien, C.E. and Jones, R.L. (2003): Early and Middle Pleistocene vegetation of the Médoc region, southwest France. *Journal of Quaternary Science* 18 (6): 557-579
- Oren Benaroya, R., Tzin, V., Tel-Or, E. and Zamski, E. (2004): Lead accumulation in the aquatic fern *Azolla filiculoides*. *Plant physiology and Biochemistry* 42 (7-8): 639-645
- Parthasarathy, R., Kardivel, R. and Kathaperumal, V. (2002): *Azolla* as a partial replacement for fish meal in broiler rations. *Indian Veterinary Journal* 79 (2): 144-146
- Reyes, O.S. and Fermin, A.C. (2003): Terrestrial leaf meals or freshwater aquatic fern as potential feed ingredients for farmed abalone *Haliotis asinina* (Linnaeus 1758). *Aquaculture Research* 34 (8): 593-599
- Rostański, K. and Sowa R., (1986-1987): Alphabetical list of the ephemerophytes of Poland. *Fragmenta Floristica et Geobotanica* 31-32(1-2): 151-203 [in Polish with English summary]
- Rune, F. and Jorgensen, H. (1997): *Andemadsbregne (Azolla Lamarck) – botanik, udbredelse og anvendelse*. *URT* 2: 59-65
- Sanyahumi, D., Duncan, J.R., Zhao, J.R., Van Hille, R. (1998): Removal of lead from solution by the non-viable biomass of the water fern *Azolla filiculoides*. *Biotechnology Letters* 20: 745-747.
- Shiny, K.J., Remani, K.N., Jalaja, T.K. and Sasidharan, V.K. (2004): Removal of chromium by two aquatic pteridophytes. *Indian Journal of Environmental Health* 46 (3): 249-251

- Shiomi, N. and Kitoh, S. (2001): Culture of *Azolla* in a pond, nutrient composition, and use as a fish feed. *Soil Science and Plant Nutrition* 47 (1): 27-34
- Stepniewska, Z., Bennicelli, R.P., Balakhnina, T.I., Szajnocha, K., Banach, A. and Wolinska, A. (2005): Potential of *Azolla caroliniana* for the removal of Pb and Cd from wastewaters. *International Agrophysics* 19 (3): 251-255
- Svenson, H.K. (1944): The New World Species of *Azolla*. *American Fern Journal* 34 (3): 69-85
- Tung, H.F. and Watanabe, I. (1983): Differential response of *Azolla-Anabaena* associations to high temperatures and minus phosphorus treatments. *New Phytologist* 93: 423-431
- Vlek, P.L.G., Eberhardt, U. and Mie Aung, M. (2002): The role of *Azolla* in lowering the pH of simulated floodwater. *Journal of Applied Botany* 76 (1-2): 1-7
- Watanabe, I. and Berja, N.S. (1983): The growth of four species of *Azolla* as affected by temperature. *Aquatic botany* 15: 175-185
- West, R.G. (1953): The occurrence of *Azolla* in British interglacial deposits. *New Phytologist* 52: 267-272
- Wolkowycki, D. (1999): *Azolla filiculoides* (Pteridophyta, Azollaceae) in Poland. *Fragmenta Floristica et Geobotanica Series Polonica* 6: 165-170 [in Polish with English summary]
- Wong Fong Sang, H.W., Vu, Van Vu, Kijne, J.W., Vu, T.T. and Planque, K. (1987): Use of *Azolla* as a test organism in a growth chamber of simple design. *Plant and Soil* 99: 219-230
- Zhao, M. and Duncan, J.R. (1998): Removal and recovery of nickel from aqueous solution and electroplating rinse effluent using *Azolla filiculoides*. *Process Biochemistry* 33 (3): 249-255
- Zhao, M., Duncan, J.R. and Sanyahumi, D. (1999a): Competitive sorption of multiple heavy metals by *Azolla filiculoides*. *Resource and Environmental Biotechnology* 2 (3): 173-183
- Zhao, M., Duncan, J.R. and Van Hille, R.P. (1999b): Removal and recovery of zinc from solution and electroplating effluent using *Azolla filiculoides*. *Water Research* 33 (6): 1516-1522

Date of creation/modification of this species fact sheet: 09-05-2007