

NOBANIS - Invasive Alien Species Fact Sheet

Homarus americanus

Authors of this factsheet: Gro I. van der Meeren¹, Josianne Støttrup², Mats Ulmestrand³, Jan Atle Knutsen¹. ¹Institute of Marine Research, PoBox 1870 Nordnes,, NO-5817 Bergen, Norway, ²Danish Institute for Fisheries Research, Kavalerigården 6, DK-2920 Charlottenlund, Denmark, ³Institute of Marine Research, P.O. Box 4, 453 21 Lysekil, Sweden.

Bibliographical reference – how to cite this fact sheet:

van der Meeren, G. , Støttrup, J., Ulmestrand, M., Knutsen, J.A. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Homarus americanus*. – From: Online Database of the North European and Baltic Network on Invasive Alien Species - NOBANIS www.nobanis.org, Date of access x/x/200x.

Species description

Scientific name: *Homarus americanus*, Milne Edwards 1837, Nephropidae, Decapoda.

Synonyms: *Astacus marinus* Say 1817; *Astacus americanus* Stebbing 1893; *Homarus mainensis* Berrill 1951

Common names: Amerikansk hummer (DK, NO, SE), American lobster (GB) Ameerika homaar (EE).



Fig. 1. *Homarus americanus*, photo by Astrid Woll, Møre Research, Ålesund.

Species identification

Clawed lobster with powerful tail, one pair of long antennae, shorter, branched antennulas, differentiated large claws and four pairs of walking legs. The colouration is usually a dark bluish green to greenish brown, with a reddish tint on body and claws, and a greenish tint on walking legs. Very dark greenish black spots on carapace and orange to whitish below. Variations in colour composition is common, but white spots near the eyes and mouth, or marbled colouration, commonly found in European lobsters (*Homarus gammarus* L.1758) are not known. The protruding rostrum between the eyes has several lateral teeth and usually on or more ventral tooth. The body length can reach 50 cm and body weight can reach up to 20 kg, although large specimens are rare (Wolff 1978). See Williams (1987) and Holthuis (1991), for full descriptions.

Native range

Northeastern American coast and waters from Cape Hatteras, Carolina in USA to Labrador, Newfoundland and Straits of Belle Isle in Canada. Depth range: from the shore to 700 m depth (See Lawton & Lavalli 1995).

Alien distribution

History of introduction and geographical spread

Two apparently American lobsters caught in Icelandic waters were brought to the Marine Research Institute in the 1960s. One was caught by a Nephropstrawl at 150-155 m depth southwest of Iceland in May 1965. The specimen was a male measuring 39 cm from rostrum to telson (total body length). The colour was greenish black. The carapace length/carapace width measurements were identical to those of *H. americanus* from Canada and USA but differed greatly from the ratio for *H. gammarus* in Europe. This was confirmed by lobster specialists both from Canada and England, although no genetical identifications were available at that time. The other lobster, a female, was caught in the early 1960s off the west coast of Iceland, but no further details on that specimen are available (Skúladóttir 1968).

The most recent documented finds in Europe were in the inner Oslofjord, Norway, November 1999, followed up by additional two landed in the same region. Five have been caught near Sandefjord, Kristiansand and Ålesund in southern and western Norway in the following three years (van der Meer *et al.* 2000). Three more were caught in 2005, west of Bergen, Norway, and positively identified as American lobsters, followed by at least four more caught by a target fishery in the same area. Some possible specimens are yet to be genetically analysed.

It is also possible that American lobsters were caught in the English Channel, but this has not been officially reported. All Atlantic, North Sea and possible Mediterranean coastal waters of Europe with native stocks of the European lobster must be considered as potential areas for introductions. Intentional implants to areas outside the range of the *Homarus* species, like the American West coast, Japan and Oceania, have not been successful.

Pathways of introduction

The introduction pathway to Norway has not been investigated, but is presumably based on release of American lobsters from live import, legal and illegal.

Unconfirmed information is that the American lobsters in the Oslofjord area was released: a) from a quayside fish-salesman, throwing out the rest of a batch imported lobsters he could not sell, b) from intentional releases by a local fisherman, illegally importing American lobsters for the purpose, c) possibly from a school, hatching, growing and possibly releasing American lobsters as a class project. The release near Kristiansand is allegedly from an illegally imported batch of live American lobsters, escaped from storage in submerged cages in the sea, while the lobsters found outside Sandefjord, Ålesund and Bergen have unknown origin. The lobsters caught in Ålesund died within few days, suggesting they might have been held by restaurants on land or cruise ships for servings, until they were discarded and released to the sea due to weak condition. All the four regions where American lobsters have been found are close to airports with international traffic, harbour with large ferry lines and visiting cruise ships and live marine crustaceans offered for sale on the market.

In the rest of Europe, a large part of the European import of live American lobsters is stored on both sides of the English Channel, in Belgium, the Netherlands, UK and France. France was also

releasing a few clutches of hatched and cultivated American lobsters in a sea ranching project in the late 1970's.

Alien status in region

All the positively identified specimens are found incidentally in Norway, while none have been observed in Germany, Denmark and Sweden (see table 1). Faroe islands, Iceland and Greenland as well as the Barent Sea are north of the climatic range of any of the *Homarus* species, although an early, not genetically certified find, was made in Icelandic waters (see above). Most of the Baltic is probably too limnic for the lobsters to survive.

Country	Not found	Not established	Rare	local	Common	Very common	Not known
Denmark	X						
Estonia	X						
European part of Russia	X						
Finland	X						
Faroe Islands	X						
Germany	X						
Greenland	X						
Iceland		X					
Latvia	X						
Lithuania	X						
Norway		X					
Poland	X						
Sweden	X						

Table 1. The frequency and establishment of *Homarus americanus*, please refer also to the information provided for this species at www.nobanis.org/search.asp. Legend for this table: **Not found** –The species is not found in the country; **Not established** - The species has not formed self-reproducing populations (but is found as a casual or incidental species); **Rare** - Few sites where it is found in the country; **Local** - Locally abundant, many individuals in some areas of the country; **Common** - Many sites in the country; **Very common** - Many sites and many individuals; **Not known** – No information was available.

Ecology

Habitat description

American lobsters are found in a wide variety in their native range. Inshore populations are found in shallow waters on mud, cobble, bedrock, peat reefs, eelgrass beds and occasionally within sandy depressions (Lawton & Lavally 1995). Offshore populations living down to 200 m depth, are found on similar substrates, as well as clay. The temperature in this area ranges from 5° C to 20° C, although the species tolerate from –1° C to 30.5° C. They can also survive shortly in low salinity, although they are usually found in areas with salinity above 25 psu.

In the invaded areas, American lobsters have been caught between 10 and 50 m depth near the coast, in habitats consisting of a mixture of bedrock, sand, mud and cobble. All have been caught in typical habitats for native crabs and lobster species, as they have been caught in lobster or crab pots.

Reproduction and lifecycle

The life cycle of the American lobster is, in many aspects, similar to the European lobster. Female lobsters mate immediately after a moult, build up the internal egg-clutch during the following year, spawn the next summer, when the eggs get fertilized during transfer and attachment to the pleopods underneath the tail. The embryonic development takes another 9-16 months, when the prelarvae are hatched. The planktonic prelarvae moult into the first larvae stage within hours, followed by three more planktonic larval stages, where the fourth is called a postlarva. The development time ranges from 22 to 103 days, dependent of the temperature. Total length of the first stage larva is on average 7.8 mm, while the postlarva reach on average 12.6 mm (Ennis 1995). The postlarvae seek to the bottom and shelter within the habitat, where they moult and become juveniles, which remain in shelter for the next two-three years. With increasing size, the juveniles will gradually move more extensively outside the shelters. They are considered to have reached the adult phase at a size of 50 mm in carapace length. This can take four years or more from hatching, dependent of temperature and habitat. Functional maturity is when they are able to mate, and produce fertilized eggs. American lobsters are known to be long-lived. Some specimens are known to be up to 100 years old and more in aquariums (Lawton & Lavalli 1975).

The habitat and food preference of the American lobster seems to be similar to the European lobster. The offshore populations of the American lobster live in burrows similar to the Norwegian lobster (*Nephrops norvegicus* L. 1758) (Chapman 1980). They also inhabit the same depth range. They might therefore be a potential space and food competitor also to this species.

The fecundity of the American lobster female is dependent on the size of the female, which can produce from a few thousand to several tens of thousands eggs per clutch. Smaller females tend to moult and spawn every second year, while larger females can produce egg-clutches two years in a row before moulting the third year (Talbot & Helluy 1995). As the larval stages have relatively short durations, where stage 3 and 4 are actively seeking into the shoreline, the dispersal is not expected to be very wide. Due to the many years it takes from mating to maturation, the population growth rates are slow, but if all life stages are established, it will be impossible to eradicate the species from invaded areas.

Dispersal and spread

It is not known if the American lobsters in Norwegian waters have produced offspring. However, one female in the Oslofjord-catches was carrying eggs, where some seemed to have been hatched immediately before being caught (van der Meeren *et al.* 2000). However, no juvenile or adolescent lobsters have been found.

Some long-distance movements has been suggested for the lobsters escaping from the holding box near Kristiansand, Vest Agder, where the distance between the most distant lobster catches was up to 30 km to each side of the escape location. As American lobsters are known to have seasonal long-distance migrations, the potential is present (Campbell & Stasko 1985; 1986).

Impact

Affected habitats and indigenous organisms

Live American lobsters are themselves a competitor to the native European lobster and possibly Norwegian lobsters, by sharing the same habitat and food preferences (Dybern 1973; Cooper & Uzmann 1980; Chapman 1980; Lawton & Lavalli 1995; Collins 1998).

Additionally, they can carry various diseases and parasites, as the Gaffkemia, a lethal bacterial blood disease, which has led to outbreak in lobster holding facilities for imported lobsters in Europe (Wiik *et al.* 1987; Mortensen 2002). Many imported American lobsters carry with them encrusting

organisms, like barnacles and polychaeta with no harmful effect on the lobsters, but with a potential for being invasive species themselves.

Being large and long-lived predators, able to manipulate their environment by digging and shuffling substrate, they might have a long-term modern effect on the ecosystem.

Being spawned only in limited seasons, and hatched as relatively large larvae, they are not expected to be prone to enter and survive in ballast-water tanks.

Genetic effects

Due to the inherent ability in both species to distinguish between the two, natural mating will not take place as long as a mate of the same species is available (van der Meeren *et al.* 2003). Cross-species mating has been achieved in laboratories when no mate choice has been offered. Such matings produce live, fast growing and vigorous offspring (F1), with traits from both species. In some cases these were reported to be sterile (Adouin & Leglise, 1972; Carlberg *et al.* 1978; Hedgecock *et al.* 1977; Talbot *et al.* 1984; Bowser & Rosemark 1981), while in one laboratory they also produced a second F2 generation (J. Kittaka, Sanriku, Japan, pers. comm.).

Human health effects

None

Economic and societal effects (positive/negative)

The European lobster stocks in the Nordic region are quite small and vulnerable, and much smaller than in historic time. Efforts are made to enhance them (Agnalt *et al.* 1999; 2004). The presence of a competitive sibling species could, if the density of the invasive species gets high, be harmful and slow the rebuilding of the native stocks. If the diseases known only in American lobsters are spread to European waters it could/will have a serious negative effect, and in the worst case, close down in the fisheries.

It is because of the low landing volume of European lobsters that live American lobsters are imported. Frozen lobsters are of inferior quality compared to live quality, and the local fisheries cannot support the demand. European countries are the most important customers of the Canadian lobster fishery.

Management approaches

Prevention methods

The present laws for Denmark, Norway and Sweden include international agreements for the prevention of biological variation (ANON 2001; 2002, 2004a; 2004b), and instructions for handling imported live animals (van der Meeren *et al.* 2004). Further, national action plans for biodiversity and international trade agreements are present, with import-laws similar in the 3 countries, requiring veterinary certificates and including a prohibition of release of lobsters in coastal waters.

Eradication, control and monitoring efforts

In Norway there is a money reward for any caught and positively identified American lobster. Most of the public aquaria and the Institute of Marine Research (IMR) are collaborating to inform the public and receive all possible specimens delivered by professional and recreational fishermen. All findings are to be analysed genetically by IMR and registered.

Improved control routines have been employed for control of imported lobsters as well as fishmongers and the fish markets. Still, it is a problem that American lobsters imported into a EU

country are regarded as “lobsters” and cannot be traced as a species in the trade statistics if sold on to a second EU country.

Information and awareness

When the first American lobsters were identified in Norway, it caught the public attention and was followed by a broad media interest. The ensuing part governmental, part media driven information campaigns have successfully educated the public about the two lobster species and their differences. This has possibly stopped further illegal import, as no new American lobsters have been identified the last two years.

However, as long as live American lobsters can be legally imported, there is a risk that someone will violate the legislations, by accident or intentionally, and release more American lobsters in Nordic waters.

Knowledge and research

Research is undertaken in several of the Nordic countries. The investigations are closely related to the putative effect the species might have on native lobsters. Themes are amongst others: dominance in shelter conflicts, mate choice (European female with one American lobster and one European lobster male), genetic analyses to distinguish between the two species, disease outbreak in Norway.

Recommendations or comments from experts and local communities

In the case of the American lobster, it is recommended to implement the first line of defence, already with its import, since it is very difficult to control once the animals have been imported. In practice our view means that all import of live American lobsters should be forbidden, both from non-EU/EØS countries and via EU/EØS countries. If this is not possible for the Nordic countries alone within the EU/EØS regulations, we suggest that an import restriction for the whole EU/EØS area should be considered. Trade within EU concerning lobsters should be registered in the import statistics at species level and it should be possible to trace the origin. Information on, and administration of, import, and handling instructions should be established and centralised within each country. This should also be coordinated between the Nordic countries, as should the registration and management of lobster fishery in common waters. This would be essential because a restriction on live lobster import may increase demand and fishing pressure on the European lobster. If it is impossible to restrict the import of live American lobsters, there is an urgent need to launch information campaigns focusing on disseminating knowledge and risks regarding the introduction of live American lobsters in native waters.

Continuous information to the public and scientific community should be maintained, as well as collecting suspected American lobster specimens for identification at the Institute of Marine Research. It is a need for development of the genetic analytic methods to get a cost- and time-effective method that also can detect hybrid lobsters.

References and other resources

Contact persons

Gro I. van der Meer (NO) (Comparative biology, Ecology)
Institute of Marine Research, PoBox 1870 Nordnes,, NO-5817 Bergen, NORWAY
Visiting adress: Austevoll Research Station, NO-5392 Storebø, NORWAY
Phone: +47 56 18 22 68, E-mail: grom@imr.no

Jan Atle Knutsen (NO) (Norwegian lobster fisheries and management)

Institute of Marine Research, PoBox 1870 Nordnes,, NO-5817 Bergen, NORWAY
Visiting adress: Flødevigen Reserach Station, NO-4817 His, NORWAY
Phone: +47 37 05 90 28, E-mail: janak@imr.no

Josianne Støttrup (DK) (Danish lobster fisheries and management)
Danish Institute for Fisheries Research, Kavalergården 6, DK-2920 Charlottenlund, DENMARK.
Phone: +45 33 96 34 29, E-mail: jgs@dfu.min.dk

Mats Ulmestrand (SE) (Swedish lobster fisheries and management)
Institute of Marine Research, P.O. Box 4, 453 21 Lysekil, SWEDEN
Phone: +46 523 18727, e-mail: mats.ulmestrand@fiskeriverket.se

Stefan Nehring (DE) AeT umweltplanung, Bismarckstrasse 19, 56068 Koblenz, GERMANY, Tel.
+49 (0)261-1330398, E-mail: nehring@aet-umweltplanung.de

Lilika Käis (EE), Estonian Ministry of Environment, Narva mnt 7a, ruum nr 329, ESTONIA, Tel.
+3726262877, E-mail: Lilika.Kais@envir.ee

Hrafnkell Eiríksson (IS) Marine Research Institute, 121 Reykjavik, ICELAND, Tel: +354 5752125,
E-mail: keli@hafro.is

Links

[Nordisk Ministeråd](#)

EU 2004. [Statistics for fisheries and international trades](#)

Lavalli, K. 2003. Shell disease and lobsters. [Commercial Fisheries News, January 2003](#)

References

Audouin, J. & Leglise, M. 1972. Premiers résultats d'expériences relatives aux possibilités d'acclimatation de homard américain *Homarus americanus* en France. ICES CM E:34.

Agnalt, A.-L.; Jørstad, K. E.; Kristiansen, T.; Nøstvold, E.; Farestveit, E.; Næss, H.; Paulsen, O.I.; Svåsand, T. 2004: Enhancing the European Lobster (*Homarus gammarus*) Stock at Kvitsøy Islands: Perspectives of Rebuilding Norwegian Stocks. In: Leber, K. M.; Kitada, S.; Blankenship, H.L.; Svåsand, T. ed. Stock enhancement and sea ranching – developments, pitfalls and opportunities. London, UK, Blackwell Publishing. Pp 415-426.

Agnalt, A.-L.; van der Meeren, G. I.; Jørstad, K. E.; Næss, H.; Farestveit, E.; Nøstvold E.; Korsøen, E.; Ydstebø, L.; Svåsand, T. 1999: Stock Enhancement of European lobster (*Homarus gammarus*); A large-scale experiment off southwestern Norway (Kvitsøy). In: Howell, B.; Moksness, E.; Svåsand, T. ed. Stock enhancement and sea ranching. London, Blackwell Scientific Press. Pp 401-419.

ANON 2001. Biologisk mangfold Sektoransvar og samordning. Miljøverndepartementet, Norge.

ANON 2002. Report of the Working group on Pathology and Diseases of Marine Organisms. ICES C.M. 2002/F:02; Ref ACME+E: 8-11.

ANON 2004a. Sveriges genomförande av Konventionen om biologisk mångfald med avseende på främmande arter og genotyper. Dnr. 2004-02-22, Naturvårdsverket, Sverige.

ANON 2004b. Handlingsplan for biologisk mangfoldighet og naturbeskyttelse I Danmark 2004-2009. Utgivet av Miljøministeriet, Danmark

Bowser, P.R. & Rosemark, R. 1981. Mortalities of cultured lobsters, *Homarus*, associated with a molt death syndrome. *Aquaculture* 23: 11-18.

- Campbell, A. & Stasko, A.B. 1985. Movements of lobsters, *Homarus americanus*, off southwestern Nova Scotia. Canadian journal for Fisheries and Aquatic Sciences 42: 229-238.
- Campbell, A. & Stasko, A.B. 1986. Movements of lobsters (*Homarus americanus*) tagged in the Bay of Fundy, Canada. Marine Biology, 92: 393-404.
- Carlberg, J.M., van Olst, J.C. & Ford, R.F. 1978. A comparison of larval and juvenile stages of the lobsters *Homarus americanus*, *Homarus gammarus* and their hybrid. In: Proceedings World Mariculture Society 9: 109-122.
- Chapman, C.J. 1980. Ecology of juvenile and adult Nephrops. I: The Biology and Management of Lobsters, vol 2 Ecology and Management, Academic Press, New York: 143-178.
- Cooper, R.J. & Uzmann, J.R. 1980. Ecology of juvenile and adult clawed lobsters *Homarus americanus*, *Homarus gammarus*, and *nephrops norvegicus*. Division of Fisheries and Oceanography Circulations (CSIRO, Australia) 7: 187-208.
- Dybern, B.I. 1973. Lobster burrows in Swedish waters. Helgoländische Wissenschaftlichen Meeresuntersuchungen, 24: 401-414.
- Collins, K.J. 1998. Habitat selection and mobility in adult lobsters. I: The European Lobster *Homarus gammarus* (L.) Proceedings From the Seminar at Kvitsøy 1995 (red.: G.I. van der Meeren & O. Soldal). Fisken og Havet, 13: 46-56.
- Hedgecock, D., Moffett, W.L., Borgeson, W. & Nelson, K. 1977. Progress and problems in lobster broodstock development. In: Proceedings World Mariculture Society 9: 497-506.
- Holthuis, L.B. 1991. Holthuis 1991. Key to species *Homarus americanus*, *H. capensis* and *H. gammarus*. FAO Species Catalogue vol. 13: Marine Lobsters of the World
- Mortensen, S. 2002. Gaffkemi i norske hummeranlegg. Et tegn på "hull" i våre kontrollrutiner? Norsk Veterinærtidsskrift 114: 471-474.
- Lawton, P. & Lavalli, K.L. 1995. Postlarval, juvenile adolescent, and adult ecology. In: Biology of the Lobster *Homarus americanus* (ed. J.R. Factor) Academic Press, San Diego: 47-88.
- Skúladóttir, U. 1968. Ný humartegund fundin við Ísland (New lobster species found in Icelandic waters). Náttúrufræðingurinn, Vol. 38, 1968, pp. 110-112.
- Talbot, P., & Helluy, S. 1995. Reproduction and embryonic development. In: Biology of the Lobster *Homarus americanus* (Ed. J.R. Factor), Academic Press San Diego, P. 177-216.
- Talbot, P., Thaler, C. & Wilson, P. 1984. Spawning, egg attachment and egg retention in captive lobsters (*Homarus americanus*). *Aquaculture* 37: 239-249.
- van der Meeren, G.I., Chandrapavan, A. og Breithaupt, T. 2003. Makevalg hos homarid hummer; kan de gjenkjenne sin egen art? Fisken og Havet nr. 1-2003: 23 pp.
- van der Meeren, G.I, Ekeli, K.O, Jørstad, K.E og Tveite, S. 2000. Americans on the wrong side – The lobster *Homarus americanus* in Norwegian waters. ICES CM 2000/U:20
- van der Meeren, G. I., Støttrup, J., Ulmestrand, M., & J.A. Knutsen 2004. Forebyggelse av ny hummerart i nordiske farvande. Sluttrapport for Arbeidsgruppen for nordisk miljø- og fiskerisamarbeid, Nordisk Ministerråd,, 6 pp.
- Wiik, R. , Egedius, E., & Goksøyr, J. 1987. Screening of Norwegian lobsters *Homarus gammarus* for the lobster pathogen *Aerococcus viridans*. Diseases in Aquatic Organisms, 3: 97-100.
- Williams, A.B. 1984. Shrimps, Lobsters , and crabs of the Atlantic Coast of the United States, Maine to Florida. Smithsonian Institution Press, Washington, D.C.
- Wolff, T. 1978. Maximum size of lobsters (*Homarus*)(Decapoda, Nephropidae). Crustaceana 34: 1-14.

Date of creation/modification of this species fact sheet: 03-06-2005