

NOBANIS - Invasive Alien Species Fact Sheet

Salmo salar

Author of this fact sheet: Peder Fiske, Norwegian Institute for Nature Research, Tungasletta 2, NO-7485 Trondheim, Norway, tel: + 47 73 80 15 22, E-mail: Peder.Fiske@nina.no

Bibliographical reference – how to cite this fact sheet:

Fiske, P. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Salmo salar*. – From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org, Date of access x/x/200x.

Species description

Scientific names: *Salmo salar* (Linnaeus, 1758), Salmonidae.

Note: this fact sheet focuses on escaped *Salmo salar*; the species as such is native to the region.

Synonyms: none

Common names: Atlantic salmon (GB), Laks (DK), Lachs (DE), Lõhe (EE), Lax (IS), lasis (LV), Atlanto lašiša (LT), Laks (NO), Lax (SE).



Fig. 1. Escaped farmed salmon (*Salmo salar*) (right) often show wavy fin rays, while wild salmon (left) has more even rays, photos by Roar A. Lund.




Fig. 2. Salmon that recently has escaped from net pens (right) often have rounded tails, while wild salmon (left) have more sharp tails, photos by Roar A. Lund.

Species identification

The appearance of Atlantic salmon changes from the juvenile freshwater stage, via a silvery stage in the ocean to a more ornamented appearance close to spawning in the rivers. Adult *Salmo salar* in the sea has silvery sides and belly, and a darker back. They normally have relatively few spots below the lateral line, a relatively slender caudal peduncle (tail root), and a v-formed tail. *Salmo salar* and sea-trout (*Salmo trutta*) may be difficult to separate, especially large individuals. Farmed salmon look similar to their wild con-specifics, but they often have worn fins with wavy fin-rays and more spots both above and below the lateral line than wild salmon. Thus, farmed salmon may at occasions look similar to sea trout. Furthermore, the growth pattern in the scales distinctively separates farmed from wild salmon (Lund and Hansen, 1991; Fiske *et al.*, 2005a).

Native range

The *Salmo salar* originally occurred in every country with rivers flowing into the North Atlantic Ocean and the Baltic Sea and is thus native to the region (Mills, 1989). Today, the species distribution has decreased in its southern range on both sides of the Atlantic. In the Baltic sea large number of reared smolts have been released to supplement local populations (for example: Saloniemi *et al.*, 2004). In Poland the last salmon population were lost from the Drawa River in the mid of the 1980s. For restoration of salmon in Polish waters Daugava salmon were used. The first salmon smolt stocking was in 1994. In 1996 the first spawners in rivers were observed. Returning salmon has been captured and used for artificial spawning (Bartel 2001). *Salmo salar* has been restocked in Latvia since 1885 (Latvijas Daba)

Alien distribution

History of introduction and geographical spread

Farming of *Salmo salar* in net pens in the sea started in 1969 in Norway (Gjedrem, 1981). In 1971 a systematic selection program started, based on broodstock salmon sampled from 40 different Norwegian rivers and one Swedish river (Gjedrem *et al.*, 1991). The production of farmed salmon in Norway has increased tremendously from 100 tonnes in 1971 to 509 554 tonnes in 2003 (Gjedrem, 1981; Hansen *et al.*, 2005). Today, *Salmo salar* is produced within (main producers: Norway, Scotland, Ireland, USA and Canada) and outside (main producers: Chile, USA and Canada) the species natural range. Farmed *Salmo salar* today by far outnumber their wild con-specifics (Gross, 1998). In the 1980s escaped farmed salmon started to be noted in catches from Norwegian rivers and sea fisheries (Moen and Gausen, 1989; Gausen and Moen, 1991), and from 1989 river and sea fisheries in Norway have been surveyed yearly for the occurrence of escaped farmed salmon (Lund *et al.*, 1991; Fiske *et al.*, 2001). Norwegian strains of salmon were introduced to Iceland in 1989 for rearing in land based facilities and later in sea-cages.

Pathways of introduction

The reported number of escaped farmed salmon from Norwegian fish farms has varied between approximately 250 000 and approximately 600 000 individuals yearly since 1994 (Norwegian fish – statistics – [on line](#)). As a comparison, the yearly population of adult salmon returning to the Norwegian coast has been estimated to have varied between approximately 500 000 and 1 000 000 individuals in the same period (Hansen *et al.*, 2005). The real number of escapees is probably higher than the reported numbers as some fish may escape without being noticed by the fish farmers. Even though many of the escapees die before they are ready to spawn, many also reach the rivers and spawn together with wild *Salmo salar* (Lura *et al.*, 1993). Progeny of escaped farmed salmon and hybrids between escaped farmed salmon and local wild salmon seem to have lower success than progeny from wild salmon (Fleming *et al.*, 1996; Fleming *et al.*, 2000; McGinnity *et*

al., 2003; McGinnity *et al.*, 2004), but still genes from the farmed salmon can enter the wild populations. This may lead to less genetic differentiation among wild populations because many of them are affected by repeated introgressions from farmed salmon (Tufto, 2001).

Alien status in region

Escaped farmed salmon have been found in catches of wild salmon in Norway (Fiske *et al.*, 2001), the British Isles (Webb *et al.*, 1993; Crozier, 1998; Milner and Evans, 2003), Iceland (Gudjonsson, 1991) and in the high seas close to the Faroe Islands (Hansen *et al.*, 1999), see also Table 1. Relatively few farm escapees are probably found in the Baltic Sea, but they can be found in Danish rivers (Niels Jepsen, pers. comm.). In Finland, fish farm escapees can occasionally be found in the River Tana forming the border between Norway and Finland (Fiske *et al.*, 2001). Note that since *Salmo salar* occurs naturally in the region, the fish farm escapees are found together with their wild con-specifics and are not as an alien species as such.

Country	Not found	Not established	Rare	Local	Common	Very common	Not known
Denmark			X				
Estonia	X						
European part of Russia							X
Finland			X				
Faroe Islands					X		
Germany	X						
Greenland			X				
Iceland				X			
Latvia				X			
Lithuania			X				
Norway						X	
Poland			X				
Sweden			X				

Table 1. The frequency and establishment of escaped *Salmo salar*, please refer also to the information provided for this species at www.nobanis.org/search.asp. Legend for this table: **Not found** –The species is not found in the country; **Not established** - The species has not formed self-reproducing populations (but is found as a casual or incidental species); **Rare** - Few sites where it is found in the country; **Local** - Locally abundant, many individuals in some areas of the country; **Common** - Many sites in the country; **Very common** - Many sites and many individuals; **Not known** – No information was available.

Ecology

Habitat description

Salmo salar juveniles are found in a wide variety of rivers draining to the sea (Elliott *et al.*, 1998). The habitat requirements varies among different life stages (Bardonnet and Baglinière, 2000), as the young will be found in faster currents and in deeper water as they grow older (Heggenes *et al.*, 1999). Where the two species are found together *Salmo salar* seem to prefer areas with faster water currents than brown trout (Heggenes *et al.*, 1999). Offspring of escaped farmed salmon will probably use similar habitats as wild salmon, and thus compete with the wild fish.

The size of adult salmon vary among rivers, and rivers seem to have to be above a certain size to carry populations of large multi-sea-winter salmon (Jonsson *et al.*, 1991). Escaped farmed salmon seems to enter the rivers later in the season than wild salmon (Fiske *et al.*, 2001).

Postsmolts and adults of *Salmo salar* are distributed over large areas in the North Atlantic Ocean during their migrations between fresh water and their feeding grounds at sea (Hansen and Jacobsen, 2000). The smolts seems to enter the sea in spring or summer when the sea surface temperature in coastal areas are above 8 °C (Hvidsten *et al.*, 1998). Their survival seems to be correlated with the distribution of relatively warm sea water in the areas they reach shortly after their entrance to the sea (Friedland *et al.*, 1998). Escaped farmed salmon occur in the same feeding grounds in the ocean as wild salmon, though more farm escapees have been found close to the Faroe Islands (Hansen *et al.*, 1993) than close to Greenland (Hansen *et al.*, 1997).

Reproduction and life cycle

Several authors have described the life cycle of the *Salmo salar* in detail (historical review in: Mills, 1989). *Salmo salar* spawn in rivers in the autumn, and the fry hatch in the spring or early summer. The young spend from 1-6 years in the river before they move to sea as smolts. In the ocean they grow fast and return to the rivers after one to five years (normally 1-3) in the sea. In larger rivers most males return after one year at sea, while most females return after two years at sea. In smaller rivers both most males and females return after one year at sea. Not all salmon die after spawning and a variable proportion spawn repeatedly, seemingly more so in populations of mostly one-sea-winter salmon than in populations of multi-sea-winter salmon (Jonsson *et al.*, 1991). Offspring of escaped farmed salmon will probably behave rather similarly to wild salmon, though they seem to incur higher mortality than wild salmon (Fleming *et al.*, 2000; McGinnity *et al.*, 2003).

Dispersal and spread

Farmed smolts that are released into the wild tend to return to the same area where they were released (Hansen and Jonsson, 1991), while adult salmon that escape tend to disperse more widely (Hansen, 2005; Hansen, 2006a; Hansen, 2006b; Skilbrei *et al.*, 2006).

Impact

Affected habitats and indigenous organisms

Wild populations of *Salmo salar* and brown trout may be affected directly by competition or interference at the spawning grounds (Lura and Sægrov, 1991; Fleming *et al.*, 1996), indirectly by gene flow from farmed to wild populations (McGinnity *et al.*, 1997; Fleming *et al.*, 2000; McGinnity *et al.*, 2004), or through the spreading of diseases or parasites (Heuch and Moe, 2001; Bjørn and Finstad, 2002).

Genetic effects

Theoretical models suggest that if high proportions of escaped farmed salmon continue to spawn together with wild salmon, the genetic variation among wild populations may diminish and the wild stocks may become genetically more similar to farmed salmon than at present (Tufto, 2001; Hindar *et al.*, 2005; Hindar *et al.*, 2006). In some rivers with high occurrences of escaped farmed salmon and weak populations of wild salmon there also appear to have been a change in the genetic profile over time (Skaala *et al.*, 2005). There are indications that the occurrence of escaped farmed salmon in rivers may have lead to increased occurrence of hybrids between *Salmo salar* and brown trout (Youngson *et al.*, 1993; Matthews *et al.*, 2000).

Human health effects

Consumption of escaped farmed fish recently treated with antibiotics may lead to the development of pathogens with resistance towards antibiotics. However, the use of antibiotics in the farming industry has decreased dramatically since the 1980s (Grave and Horsberg, 2005).

Economic and societal effects (positive/negative)

The salmon farming industry is one of Norway's largest export industries, providing work for people in remote areas of the country. River owners and anglers in several countries (Norway, Island) have raised concerns that the image of the highly priced angling fisheries for salmon may be harmed if high numbers of escaped farmed salmon enter the rivers. Also in other countries angling of *salmo salar* is very important – as an example the annual catch of this species is 350 - 600 tons in Latvia (Latvijas Daba).

Management approaches

Prevention methods

The fish farming industry has increased their efforts to prevent escapes from happening. This includes new technical standards for fish farms, overviews of the most common causes of escapes to learn from previous mistakes, and increased awareness of the problem by the personnel working at the fish farms (Valland, 2005).

Eradication, control and monitoring efforts

Shortly after reported escapes, it is common practice to allow fishing with nets in the sea close to the site of the escape. Furthermore, fishing with nets and bag-nets in the sea has also been allowed in the autumn in many counties in Norway (e.g. Syvertsen and Vatne, 2000; Fiske, 2004). This is a fishery mainly targeting escaped farmed salmon since most wild salmon are either in the rivers or at the high seas at that time. Fishing in the outlets of rivers in the autumn with rods or bag-nets has also been used to reduce the number of farmed salmon entering rivers (Fiske *et al.*, 2005b). A monitoring program to assess the proportion of escaped farmed salmon in catches and in spawning populations has been running in Norway since 1989 (Fiske *et al.*, 2001). In Island 10 % of the salmon in sea-cages have to be tagged. The usual tagging method is microtags with adipose fin clips as an external mark. Estonia has been running a program for monitoring the ratio between wild and ranched salmon

Information and awareness

The Directorate for Nature Management in Norway, Norwegian Institute for Nature Research and the Institute of Marine Research have all produced information brochures about escaped farmed salmon. In Iceland the use of tags for identifying salmon from sea-cages is advertised and a lottery is used for enhancement of tag returns.

Knowledge and research

There has been much research about escaped farmed salmon, the latest update about both escapees and other impacts of fish farming was given at a symposium in Bergen in October 2005 (Anon, 2005). Information presented at this symposium is published in a special issue of ICES Journal of Marine Science (Vol. 63, no 7, August 2006). Two similar symposia have previously been arranged in Loen, Norway (1991) and in Bath, Great Britain (1997).

Recommendations or comments from experts and local communities

Even though the fish farming industry has increased its awareness of escapees, it is important that the industry continue to work with improving their standards in handling of fish in order to keep the fish inside the net pens. Little can be done once the fish has escaped so the most important efforts are those that are done to prevent salmon from escaping.

References and other resources

Contact persons

Peder Fiske, Norwegian Institute for Nature Research, Tungasletta 2, NO-7485 Trondheim, Norway, Tel.: + 47 73 80 15 22, E-mail: Peder.Fiske@nina.no

Gudni Gudbergsson (IS), Institute of Freshwater Fisheries, Vagnhofdi 7, 110 Reykjavik, Iceland. Tel: +354 5676400. Fax: +354 5676420, E-mail: gudni.gudbergsson@veidimal.is

Niels Jepsen (DK) Afd. for Ferskvandsfiskeri (FFI), Danmarks Fiskeriundersøgelser, Vejløvej 39, DK-8600 Silkeborg, Denmark, Tel.: + 45 33963131, E-mail: nj@difres.dk

Michael Møller Hansen (DK) Afd. for Ferskvandsfiskeri (FFI), Danmarks Fiskeriundersøgelser, Vejløvej 39, DK-8600 Silkeborg, Denmark, Tel.: +45 33963145, E-mail: mmh@difres.dk

Dr. Stefan Nehring (DE) AeT umweltplanung, Bismarckstrasse 19, 56068 Koblenz, Germany, Tel.: +49 (0)261 - 13303-98, E-mail: nehring@aet-umweltplanung.de

Christina Birnbaum (EE), Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Kreutzwaldi 64, Tartu, Estonia, E-mail: chbirnbaum@yahoo.com

Jaakko Erkinaro (FIN), Finnish Game and Fisheries Research Institute, Tutkijantie 2, FIN-90570 Oulu, Finland, Tel: +358 205751871, E-mail: jaakko.erkinaro@rktl.fi

Viktoras Didžiulis (LT) Coastal Research and Planning Institute, Klaipeda University, Lithuania, H.Manto - 84, LT-91210, Klaipėda, Phone: +370 46 380905, E-mail: viktoras@ekoinf.net

Nora Kabuce, Latvian Environment, geology and meteorology agency, Maskavas Str. 165, Riga, LV-1019, Latvia. Tel.: +371 7032030, E-mail: nora.kabuce@lvgma.gov.lv

Melanie Josefsson (SE), Swedish Environmental Protection Agency, SE106 48 Stockholm, Sweden, Tel: +46 18 673148, E-mail: Melanie.Josefsson@snv.slu.se

Ryszard Bartel (PL) Iland Fisheries Institute, Departmen of Migratory Fish, Reduta Żbik str.5., 80-761 Gdansk, Poland, E-mail: gdansk@infish.com.pl

Links

[Norwegian Institute for Nature Research](#)

[Institute of Marine Research](#)

[Directorate for Nature Management](#)

[Statistics Norway](#)

[Directorate for Fisheries Management - statistics](#)

[Institute of Freshwater Fisheries](#) (Icelandic)

References

- Anon. 2005. Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions - Programme & Abstracts Bergen.
- Bardonnet, A., and Baglinière, J. L. 2000. Freshwater habitat of Atlantic salmon (*Salmo salar*). Canadian Journal of Fisheries and Aquatic Sciences, 57:497-506.
- Bartel R. 2001. Return of salmon back to Polish waters. Ecohydrol. & Hydrobiol., 1, 3, 377-392.
- Bjørn, P. A., and Finstad, B. 2002. Salmon lice, *Lepeophtheirus salmonis* (Krøyer), infestation in sympatric populations of Arctic char, *Salvelinus alpinus* (L.), and sea trout, *Salmo trutta* (L.), in areas near and distant from salmon farms. ICES Journal of Marine Science, 59:131-139.
- Crozier, W. W. 1998. Incidence of escaped farmed salmon, *Salmo salar* L., in commercial salmon catches and fresh water in Northern Ireland. Fisheries Management and Ecology, 5:23-29.
- Elliott, S. R., Coe, T. A., Helfield, J. M., and Naiman, R. J. 1998. Spatial variation in environmental characteristics of Atlantic salmon (*Salmo salar*) rivers. Canadian Journal of Fisheries and Aquatic Sciences, 55 (Suppl. 1):267-280.
- Fiske, P. 2004. Sammenstilling av data fra undersøkelser om fiske etter rømt oppdrettslaks (DN's ref. nr. 03040045). Notat, til DN 01.07.2004:1-11.
- Fiske, P., Lund, R. A., and Hansen, L. P. 2005a. Identifying fish farm escapees. In: Stock Identification Methods pp. 659-680. Ed. by S. X. Cadrin, K. D. Friedland and J. R. Waldman) Elsevier Academic Press. Amsterdam:
- Fiske, P., Lund, R. A., Østborg, G., Heggberget, T. G., and Thorstad, E. B. 2005b. Rømt oppdrettslaks i Salvassdraget i 2004 - kommer fiskene fra en eller flere rømmingsepisoder. NINA Minirapport, 103:1-9.
- Fiske, P., Lund, R. A., Østborg, G. M., and Fløystad, L. 2001. Rømt oppdrettslaks i sjø- og elvefisket i årene 1989-2000. NINA Oppdragsmelding, 704:1-26.
- Fleming, I. A., Hindar, K., Mjølnerød, I. B., Jonsson, B., Balstad, T., and Lamberg, A. 2000. Lifetime success and interactions of farmed salmon invading a native population. Proceedings of the Royal Society of London B, 267:1517-1523.
- Fleming, I. A., Jonsson, B., Gross, M. R., and Lamberg, A. 1996. An experimental study of the reproductive behaviour and success of farmed and wild Atlantic salmon (*Salmo salar*). J. Appl. Ecol., 33:893-905.
- Friedland, K. D., Hansen, L. P., and Dunkley, D. A. 1998. Marine temperatures experienced by postsmolts and the survival of Atlantic salmon, *Salmo salar* L., in the North Sea area. Fisheries Oceanography, 7:22-34.
- Gausen, D., and Moen, V. 1991. Large-scale escapes of farmed Atlantic salmon (*Salmo salar*) into Norwegian rivers threaten natural populations. Canadian Journal of Fisheries and Aquatic Sciences, 48:426-428.
- Gjedrem, T. 1981. Oppdrett av laks og aure. Landbruksforlaget, Oslo
- Gjedrem, T., Gjøen, H. M., and Gjerde, B. 1991. Genetic origin of Norwegian farmed Atlantic salmon. Aquaculture, 98:41-50.
- Grave, K., and Horsberg, T. E. 2005. Trends in the usage of drugs in Norwegian aquaculture. In: *Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions*, pp. 28. Bergen 18-21 October 2005: ICES/NASCO.
- Gross, M. R. 1998. One species with two biologies: Atlantic salmon (*Salmo salar*) in the wild and in aquaculture. Canadian Journal of Fisheries and Aquatic Sciences, 55 (Supplement 1):131-144.
- Gudjonsson, S. 1991. Occurrence of reared salmon in natural salmon rivers in Iceland. Aquaculture, 98:133-142.
- Hansen, L. P. 2005. Migration and survival of farmed Atlantic salmon (*Salmo salar* L.) released from two Norwegian fish farms. In: *Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions*, pp. 16-17. Bergen 18-21 October 2005: ICES/NASCO.
- Hansen, L. P. 2006a. Migration and survival of farmed Atlantic salmon (*Salmo salar* L.) released from two Norwegian fish farms. ICES Journal of Marine Science, 63:1211-1217.
- Hansen, L. P. 2006b. Vandrings og spredning av rømt oppdrettslaks. NINA Rapport, 162:1-21.
- Hansen, L. P., Fiske, P., Holm, M., Jensen, A. J., and Sægrov, H. 2005. Bestandsstatus for laks i Norge 2004. Rapport fra arbeidsgruppe. Utredning for DN, 2005-4:1-44.
- Hansen, L. P., and Jacobsen, J. A. 2000. Distribution and migration of Atlantic salmon *Salmo salar* L., in the sea. In: *The Ocean Life of Atlantic Salmon* pp. 75-87. Ed. by D. Mills) Fishing News Books. Oxford:
- Hansen, L. P., Jacobsen, J. A., and Lund, R. A. 1993. High numbers of farmed Atlantic salmon, *Salmo salar* L., observed in oceanic waters north of the Faroe Islands. Aquaculture and Fisheries Management, 24:777-781.
- Hansen, L. P., Jacobsen, J. A., and Lund, R. A. 1999. The incidence of escaped farmed Atlantic salmon, *Salmo salar* L., in the Faroese fishery and estimates of catches of wild salmon. ICES Journal of Marine Science, 56:200-206.
- Hansen, L. P., and Jonsson, B. 1991. The effect of timing of Atlantic salmon smolt and post-smolt release on the distribution of adult return. Aquaculture, 98:61-67.
- Hansen, L. P., Reddin, D. G., and Lund, R. A. 1997. The incidence of reared Atlantic salmon (*Salmo salar* L.) of fish farm origin at West Greenland. ICES Journal of Marine Sciences, 54:152-155.

- Heggenes, J., Bagliniere, J. L., and Cunjak, R. A. 1999. Spatial niche variability for young Atlantic salmon (*Salmo salar*) and brown trout (*S. trutta*) in heterogeneous streams. *Ecology of Freshwater Fish*, 8:1-21.
- Heuch, P. A., and Moe, T. A. 2001. A model of salmon louse production in Norway: effects of increasing salmon production and public management measures. *Diseases of Aquatic Organisms*, 45:145-152.
- Hindar, K., Fleming, I. A., McGinnity, P., and Diserud, O. 2005. Genetic and ecological interactions between wild and cultured diadromous fish. In: *Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions*, pp. 13. Bergen 18-21 October 2005: ICES/NASCO.
- Hindar, K., Fleming, I. A., McGinnity, P., and Diserud, O. 2006. The genetic and ecological effects of salmon farming on wild salmon: modelling from experimental results. *ICES Journal of Marine Science*, 63:1234-1247.
- Hvidsten, N. A., Heggberget, T. G., and Jensen, A. J. 1998. Sea water temperature at Atlantic salmon smolt entrance. *Nordic J. Freshw. Res.*, 74:79-86.
- Jonsson, N., Hansen, L. P., and Jonsson, B. 1991. Variation in age, size and repeat spawning of adult Atlantic salmon in relation to river discharge. *Journal of Animal Ecology*, 60:937-947.
- "Latvijas Daba" (Nature of Latvia) - online encyclopaedia ([link](#))
- Lund, R. A., and Hansen, L. P. 1991. Identification of wild and reared Atlantic salmon, *Salmo salar* L., using scale characters. *Aquaculture and Fisheries Management*, 22:499-508.
- Lund, R. A., Økland, F., and Hansen, L. P. 1991. Farmed Atlantic salmon (*Salmo salar*) in fisheries and rivers in Norway. *Aquaculture*, 98:143-150.
- Lura, H., Barlaup, B. T., and Saegrov, H. 1993. Spawning behaviour of a farmed escaped female atlantic salmon (*Salmo salar*). *Journal of Fish Biology*, 42:311-313.
- Lura, H., and Sægrov, H. 1991. Documentation of successful spawning of escaped female Atlantic salmon, *Salmo salar*, in Norwegian rivers. *Aquaculture*, 98:151-159.
- Matthews, M. A., Poole, W. R., Thompson, C. E., McKillen, J., Ferguson, A., Hindar, K., and Wheelan, K. F. 2000. Incidence of hybridization between Atlantic salmon, *Salmo salar* L., and brown trout, *Salmo trutta* L., in Ireland. *Fisheries Management and Ecology*, 7:337-347.
- McGinnity, P., Prodöhl, P., Ferguson, A., Hynes, R., Ó Maoiléidigh, N., Baker, N., Cotter, D., O'Hea, B., Cooke, D., Rogan, G., Taggart, J., and Cross, T. 2003. Fitness reduction and potential extinction of wild populations of Atlantic salmon, *Salmo salar*, as a result of interactions with escaped farm salmon. *Proceedings of the Royal Society of London B*, 270:2443-2450.
- McGinnity, P., Prodöhl, P., Ó Maoiléidigh, N., Hynes, R., Cotter, D., Baker, N., O'Hea, B., and Ferguson, A. 2004. Differential lifetime success and performance of native and non-native Atlantic salmon examined under communal natural conditions. *Journal of Fish Biology*, 65 (Supplement A):173-187.
- McGinnity, P., Stone, C., Taggart, J. B., Cooke, D., Cotter, D., Hynes, R., McCamley, C., Cross, T., and Ferguson, A. 1997. Genetic impact of escaped farmed Atlantic salmon (*Salmo salar* L.) on native populations: use of DNA profiling to assess freshwater performance of wild, farmed, and hybrid progeny in a natural river environment. *ICES Journal of Marine Science*, 54:998-1008.
- Mills, D. 1989. *Ecology and management of Atlantic salmon*. Chapman & Hall, London
- Milner, N. J., and Evans, R. 2003. The incidence of escaped Irish farmed salmon in English and Welsh rivers. *Fisheries Management and Ecology*, 10:403-406.
- Moen, V., and Gausen, D. 1989. Rømt oppdrettsfisk i vassdrag 1988. DN-Rapport-3-1989:1-32.
- Saloniemi, I., Jokikokko, E., Kallio-Nyberg, I., Jutila, E., and Pasanen, P. 2004. Survival of reared and wild Atlantic salmon smolts: size matters moire in bad years. *ICES Journal of Marine Science*, 61:782-787.
- Skilbrei, O., Holst, J. C., and Holm, M. 2006. Oppsummering av kunnskapsstatus innen rømming av oppdrettslaks - Tiltak for gjenfangst etter rømming. Havforskningsinstituttet, Rapport i trykk:1-20.
- Skaala, Ø., Wennevik, V., and Glover, K. 2005. Temporal genetic stability in Atlantic salmon populations. In: *Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions*, pp. 19. Bergen 18-21 October 2005: ICES/NASCO.
- Syvvertsen, H. A., and Vatne, T. 2000. Forsøksfiske etter rømt oppdrettslaks i Nordland. Fylkesmannen i Nordland, Miljøvernnavdelingen, Rapport februar 2000:1-17.
- Tufto, J. 2001. Effects of releasing maladapted individuals: a demographic-evolutionary model. *American Naturalist*, 158:331-340.
- Valland, A. 2005. The causes and scale of escapes from salmon farming. In: *Interactions between aquaculture and wild stocks of Atlantic salmon and other diadromous fish species: science and management, challenges and solutions*, pp. 15. Bergen 18-21 October 2005: ICES/NASCO.
- Webb, J. H., Youngson, A. F., Thompson, C. E., Hay, D. W., Donaghy, M. J., and McLaren, I. S. 1993. Spawning of escaped farmed Atlantic salmon, *Salmo salar* L., in western and northern Scottish rivers: egg deposition by females. *Aquaculture and Fisheries Management*, 24:663-670.

Youngson, A. F., Webb, J. H., Thompson, C. E., and Knox, D. 1993. Spawning of escaped farmed Atlantic salmon (*Salmo salar*): hybridization of females with brown trout (*Salmo trutta*). Canadian Journal of Fisheries and Aquatic Sciences, 50:1986-1990.

Date of creation/modification of this species fact sheet: 03-11-2006