

# NOBANIS - Marine invasive species in Nordic waters - Fact Sheet

## *Balanus improvisus*

---

**Author of this species fact sheet:** Kathe R. Jensen, Zoological Museum, Natural History Museum of Denmark, Universitetsparken 15, 2100 København Ø, Denmark. Phone: +45 353-21083, E-mail: [krjensen@snm.ku.dk](mailto:krjensen@snm.ku.dk)

**Bibliographical reference – how to cite this fact sheet:**

Jensen, Kathe R. (2010): NOBANIS – Invasive Alien Species Fact Sheet – *Balanus improvisus* – From: Identification key to marine invasive species in Nordic waters – NOBANIS [www.nobanis.org](http://www.nobanis.org), Date of access x/x/201x.

## Species description

### Species name

*Balanus improvisus*, Darwin, 1854 – Bay barnacle (an acorn barnacle)

### Synonyms

*B. ovularis*; *Amphibalanus improvisus* (Darwin, 1854)

### Common names

Brakvandsrur (DK); Slät havstulpan, Brackvattenlevande havstulpan (SE); Brakkvannsrur (NO); Merirokko (FI); Brakwaterpok (NL); Bay barnacle, acorn barnacle (all sessile species) (UK, USA); Ostsee-Seepocke, Brackwasser-Seepocke (DE); Pškla bałtycka (PL); Tavaline tõruvähk (EE); Jura zile (LV); Juros gile (LT); Morskoj zhelud (RU)

### Identification

*Balanus improvisus* has 6 smooth shell plates surrounding the body. It reaches a maximum size of 20 mm in diameter, but usually is less than 10 mm. The “mouth plates”, called scutum and tergum, form a diamond-shaped center. The most characteristic feature is the calcareous base with radial pattern. This base remains on the substrate after removal of the animal. There are several other species of barnacles in Nordic waters, but only two have smooth shells, *B. improvisus* and *B. crenatus* Bruguière, 1789, and the latter does not have the radial pattern on the base plate. In areas with more pronounced tides there are also species of *Chthamalus*. In Danish waters there is another introduced barnacle, [Elminius modestus \(see separate fact-sheet\)](#), and on the Swedish west coast. *B. amphitrite* Darwin, 1854, has been found (Främmande Arter, 2006).



*Balanus improvisus* on *Anodonta cygnea* from Poland.



*Balanus improvisus*, 1 specimen separated from above with the 6 smooth plates clearly visible.

For more information about *Balanus crenatus* see:

<http://www.marlin.ac.uk/speciesfullreview.php?speciesID=2718>

For information about *Balanus amphitrite* and other barnacles found introduced close to Nordic waters, see [Introduction to barnacles](#)

For more information about the native *Semibalanus balanoides* (Linnaeus, 1767) see:

<http://www.marlin.ac.uk/speciesfullreview.php?speciesID=4328>

For more information about *Chthamalus stellatus* (Poli, 1795) see:

<http://www.marlin.ac.uk/speciesfullreview.php?speciesID=2982>

and for *Chthamalus montagui* Southward, 1976 see:

<http://www.marlin.ac.uk/speciesfullreview.php?speciesID=2981>

### **Taxonomic note**

Recently, there have been some genus name changes in the barnacles, and *B. improvisus* has been transferred to the genus *Amphibalanus* Pitombo, 2004. This genus has only been described in 2004, and apparently the name change has not been widely accepted (Clare & Høeg, 2008). Since almost all of the literature on ecology and distribution of this species have used the genus name *Balanus*, this has also been used in this fact-sheet. *Balanus* is also the genus name used by the database World Register of Marine Species (WoRMS; <http://www.marinespecies.org/>).

## **Distribution**

### **Native area**

Most likely this species originates on the Atlantic coast of America, though it is uncertain whether North or South America is the source area. In some European countries, e.g. the United Kingdom, this species is considered native. Also, there are records of fossil specimens from the Pliocene in Spain (Kerckhof, 2002), and thus *B. improvisus* may have a history similar to that of *Mya arenaria* becoming extinct in Northeast Atlantic waters during the last glaciation and then re-introduced by human activities as the earliest transatlantic voyages took place. There is little genetic variation among populations in the Baltic; the U.K. and east coast of the USA (Furman, 1990).

### **Introduced area**

This species has been found in European waters for 200 years or more and must be considered fully naturalized in Nordic waters, including the Baltic Sea. There is some discussion of the earliest records. Wolff (2005) mentions that it was found in the Netherlands in 1827, and Kerckhof & Cattrijsse (2001) found it in archaeological remains from the 17th century, but it was absent from a 9th century site in Belgium (Kerckhof, 2002). Darwin (1854), in his original description of the species, mentions that it was only found a few places in England and one locality in Scotland. At the present time its distribution remains restricted to estuaries (Furman et al., 1990). The first record from the German North Sea coast is from 1858 (Gollasch & Nehring, 2006). The first record from the Atlantic coast of France is from 1872 (Gouletquer et al., 2002). The earliest record from Norway seems to be from 1900 when *B. improvisus* was found in Oslofjord, and until 1969 it was only found in this fjord. Later it has been found in several localities between Oslofjord and Stavanger, but not north of this (Sneli, 1972). It may have spread somewhat further north in recent years (Brattegard & Holthe, 1997; Hopkins, 2001). The first Swedish record is from the west coast

in 1895 (Gislén, 1950). In Ireland it is only estimated to have arrived before 1950 (Minchin, 2007).

It is more certain that the introduction to the Baltic Sea has been through human activities. Darwin had borrowed the collection of Cirripedia from the Zoological Museum in Copenhagen when he was writing his monograph (Darwin, 1854), and he does not mention *B. improvisus* from this collection. The first record of *B. improvisus* is from 1844, when it was found at Kaliningrad, then Königsberg (Leppäkoski & Olenin, 2000). The first Danish record seems to be from 1880 in the harbor of Copenhagen (Krüger, 1927; Jensen & Knudsen, 2005). It was found on the Swedish Baltic coast in 1919 (Gislén, 1950). The first record from Finland appears to be 1868 near Turku (Leppäkoski & Olenin, 2000). Presently it occurs in the Baltic up to 64° N (Leppäkoski & Olenin, 2000).

*Balanus improvisus* has also been introduced to the Black and Caspian Seas. It arrived in the Caspian Sea in 1955 (Grigorovich et al., 2003), presumably through the Volga-Don Canal (Zvyagintsev, 2003). The first record from the Black Sea dates back to 1844 (Gomoiu et al., 2002). There are no recent records from the Mediterranean, and it is unknown whether *B. improvisus* is established here (Zenetos et al., 2005). The first record may be from 1972 (Streftaris et al., 2005)

*Balanus improvisus* has also been introduced to the west coast of North America, where it was first found in 1853 in San Francisco Bay (Carlton & Zullo, 1969). Presently it is found in California, Oregon and the state of Washington (Wonham & Carlton, 2005).

The occurrence of *B. improvisus* in Japan is most likely due to human introduction. It was first recorded in 1962, and it has been spreading slowly since that time (Iwasaki, 2006). In the Russian part of the Sea of Japan *B. improvisus* is considered established. The first record was in 1969 when it was found on hydrotechnical installations (Zvyagintsev, 2003; Ovsyannikova, 2008).

Along with several other species of Cirripedia *B. improvisus* was found on an oil platform, which had been transported from Japan to New Zealand (Foster & Willan, 1979). There are also some old records of this species from ships' hulls in Australian ports, but there are no records of its having become established in these waters.

### **Vector**

Almost certainly hull fouling, though ballast water cannot be excluded for more recent introductions. As it also often grows on the shells of molluscs and large crustaceans, it may also be transferred with live commercial shellfish, and this is considered the most likely vector for the introduction to the Pacific coast of the USA (Carlton & Zullo, 1969).

## **Ecology**

*Balanus improvisus* attaches to almost any hard substrate available, such as rocks, jetties, ships' hulls, bivalve shells, macroalgae, etc. In Florida it has been found on a brackish water turtle, the diamond-backed terrapin (Ross & Jackson, 1972). In the introduced areas it will attach to the shells of local bivalves; in the Baltic it has been found on *Mytilus edulis* (Laihonen & Furman, 1986), [\*Mya arenaria\*](#) Linnaeus, 1758 (Olszewska, 2000) and even on the freshwater bivalve *Anodonta cygnea* (own observation), and in the Caspian Sea it uses an endemic bivalve *Didacna* sp. (Riedel et al.,

2006). In the Sea of Japan it settles on the native *Corbicula japonica*, which may live in freshwater, and on *Crassostrea gigas*, but also on seagrass and macroalgae (Ovsyannikova, 2008). The settling stage, the cypris larva, is probably the most important because after settling the barnacle is unable to move for the rest of its life. *B. improvisus* occurs intertidally and in shallow subtidal waters, but in the Baltic it can be found at depths of more than 50 m (Leppäkoski & Olenin, 2000). It may occur in densities of several thousand per m<sup>2</sup> (Leppäkoski, 1999; Ovsyannikova, 2008).

Like all barnacles, *B. improvisus* is a suspension feeder, creating a feeding current by pumping movements of the opercular plates and regular beating of the “cirri”, modified thoracic legs, and filtering edible particles with the setae on these cirri. The current speed and the strength of the beating of the cirri can be adjusted to the concentration and size of food particles (Crisp & Southward, 1961; Rainbow, 1984). In laboratory experiments they feed on phytoplankton, but had slower body growth than in the field (Costlow & Bookhout, 1953, 1957). When settling on *Mytilus* they prefer the posterior end of the shell, near the in- and exhalent openings of the mussel, so they can benefit from the feeding current of the mussel (Laihonen & Furman, 1986).

Growth is very fast; in three weeks a newly metamorphosed individual may have a diameter of 5 mm, and along the Swedish west coast the species may have three generations in one summer season (Stephensen, 1933). At Tvärminne in Finland they grew from 3 to a maximum of 16 mm in 10 weeks (Laihonen & Furman, 1986). *B. improvisus* molts every 2-4 days at 20° C (Costlow & Bookhout, 1953). The shell plates grow continuously as the body gets bigger. Molting can be induced and inhibited by the same hormones as in decapod crustaceans (Davis & Costlow, 1974).

*Balanus improvisus* is highly euryhaline, living in almost freshwater in some localities in the native area, La Plata River (Darwin, 1854), and the introduced area, Baltic Sea lagoons (Poulsen, 1935; Leppäkoski & Olenin, 2000). There is some disagreement about the capacity of this species to osmoregulate at low salinities (Foster, 1970; Fyhn, 1976; Rainbow, 1984). It is also rather eurythermal, tolerating temperatures as low as -2° C and up to 35° C (Southward, 1957). Furthermore it is tolerant of low oxygen concentration, eutrophication and pollution (Poulsen, 1935; Leppäkoski & Olenin, 2000; Ovsyannikova, 2008).

On the east coast of North America the flatworm *Stylochus ellipticus* Girard is a dominant predator of *B. improvisus* (Branscomb, 1976). When epibiotic on blue mussels, *B. improvisus* may be eaten with the mussel by the crab *Carcinus maenas* (Wahl et al., 1997).

## **Reproduction**

Like all barnacles, *B. improvisus* is hermaphroditic, but requires cross-fertilization, though self-fertilization has been described (Furman & Yule, 1990). Prior to copulation the barnacle acting as male briefly stops pumping water and beating the cirri, and the extremely long penis is extended into the mantle cavity of the recipient barnacle (Crisp & Southward, 1961). Egg size is about 180µm, and contrary to other species, there is little geographic variation in this size (Barnes & Barnes, 1965). *B. improvisus* may produce 1000 to 10,000 eggs per season (Costlow & Bookhout, 1957). Embryos are brooded in an ovisac inside the mantle cavity. Development to hatching takes about 21 days at 18° C (Furman & Yule, 1990).

Larvae hatch as nauplii, and a new brood can be released every 5-6 days (Gamfeldt et al., 2005). There are six nauplius stages of which the first may be non-feeding, the others feeding in the plankton, and a non-feeding cypris stage (Barnes & Barnes, 1965; Zega et al., 2007). Nauplius

larvae show strongly positive phototaxis, which decreases in the last nauplius stage (Lang et al., 1979). Both eggs and nauplii of *B. improvisus* are smaller than those of most other barnacle species (Barnes & Barnes, 1965; Ross et al., 2003). Development through the six nauplius stages takes about one week in the laboratory (O'Connor & Richardson, 1996; Dahlström et al., 2000), but this depends on temperature. Metamorphosis has been described and illustrated in great detail (Doochin, 1951). Cypris larvae are most prone to settling when they are 3-4 days old (Sjögren et al., 2008). Settlement increases in the presence of extract from adult conspecifics, and significantly more cyprids settled at 5 and 10 ppt than at other salinities between 2 and 35 ppt (Dineen & Hines, 1992). Also, the cyprids prefer rough surfaces over smooth ones, and preferentially settle in depressions (Rainbow, 1984).

## Impacts

Like other barnacles, *B. improvisus* is part of the fouling community, attaching to ships' hulls and other man-made structures. It may form dense layers on the surface of these structures, inhibiting water flow, attracting associate fauna and producing organic debris (Leppäkoski, 1999). Hence many studies have been made on factors controlling settling (Dineen & Hines, 1992; O'Connor & Richardson, 1996; Gamfeldt et al., 2005; Zega et al., 2007) and developing antifouling agents. Larvae of *B. improvisus* are used on a regular basis in toxicity tests, especially in connection with antifouling substances (Berntsson et al., 2000; Dahlström et al., 2000; Sjögren et al., 2004, 2008; Toth & Lindeborg, 2008). Neurotransmitters play an important role in barnacle settling, but different neurotransmitters may be necessary for different species. For *B. improvisus* dopamine stimulates settling whereas in *B. amphitrite* serotonin is the stimulating agent (Zega et al., 2007). In the Baltic where *B. improvisus* is the only barnacle the fouling impacts may be considerable, e.g. the costs of cleaning pipes of power plants, etc. (Leppäkoski & Olenin, 2000). However, in other areas the same impacts are done by native barnacles.

*Balanus improvisus* is a strong competitor for space (Dürr & Wahl, 2004; Gamfeldt et al., 2005), but it does not have a negative effect on community diversity in the Baltic (Dürr & Wahl, 2004).

*Balanus improvisus* has been proposed as a biomonitor for heavy metals (Rainbow et al., 2000, 2002, 2004). However, it usually does not live longer than two years and thus has limited value for long-term studies.

## References

- Barnes, H. and Barnes, M. 1965. Egg size, nauplius size, and their variation with local, geographical, and specific factors in some common cirripedes. *Journal of Animal Ecology* 34(2): 391-402.
- Berntsson, K.M., Andreasson, H., Jonsson, P.R., Larsson, L., Ring, K., Petronis, S. and Gatenholm, P. 2000. Reduction of barnacle recruitment on micro-textured surfaces: analysis of effective topographic characteristics and evaluation of skin friction. *Biofouling* 16(2-4): 245-261.
- Bousfield, E.L. 1954. The distribution and spawning seasons of barnacles on the Atlantic coast of Canada. *Bulletin of the National Museum of Canada* 132: 112-154.

- Branscomb, E.S. 1976. Proximate causes of mortality determining the distribution and abundance of the barnacle *Balanus improvisus* Darwin in Chesapeake Bay. *Chesapeake Science* 17(4): 281-288.
- Brattegard, T. & T. Holthe (Eds.) 1997. Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue. Preliminary edition. DN 1997-1. Directorate for Nature Management, Trondheim, Norway, 409pp.
- Brattström, H. 1958. Undersökningar över Öresund XXXVII. Cirripedia and Ascothoracica from the Sound and southeastern Kattegat. *Lunds Universitets Årsskrift N.F. Avd. 2*, 54(6): 1-26.
- Carlton, J.T. and Zullo, V.A. 1969. Early records of the barnacle *Balanus improvisus* Darwin from the Pacific coast of North America. *Occasional Papers of the California Academy of Sciences* 75: 1-6.
- Clare, A.S. and Høeg, J.T. 2008. *Balanus amphitrite* or *Amphibalanus amphitrite*? A note on barnacle nomenclature. *Biofouling* 24(1): 55-57.
- Costlow, J.D., Jr. 1956. Shell development in *Balanus improvisus* Darwin. *Journal of Morphology* 99: 359-415.
- Costlow, J.D., Jr. and Bookout, C.G. 1953. Moulting and growth in *Balanus improvisus*. *Biological Bulletin* 105(3): 420-433.
- Costlow, J.D., Jr. and Bookout, C.G. 1957. Body growth versus shell growth in *Balanus improvisus*. *Biological Bulletin* 113: 224-232.
- Crisp, D.J. and Southward, A.J. 1961. Different types of cirral activity of barnacles. *Philosophical Transactions of the Royal Society B* 243: 271-308.
- Dahlström, M., Mårtensson, L.G.E., Jonsson, P.R., Arnebrant, T. and Elwing, H. 2000. Surface active adrenoceptor compounds prevent the settlement of cyprid larvae of *Balanus improvisus*. *Biofouling* 16(2-4): 191-203.
- DAISIE, 2006. *Balanus improvisus*. (Fact-sheet available on [http://www.europe-aliens.org/pdf/Balanus\\_improvisus.pdf](http://www.europe-aliens.org/pdf/Balanus_improvisus.pdf), accessed on 15 March 2010).
- Darwin, C. 1854. A monograph of the subclass Cirripedia with figures of all the species. The Balanidae (or sessile cirripedes); the Verrucidae, etc. Ray Society, London.
- Davis, C.W. and Costlow, J.D. 1974. Evidence for a molt inhibiting hormone in the barnacle *Balanus improvisus* (Crustacea, Cirripedia). *Journal of Comparative Physiology* 93: 85-91.
- Dineen, J.F., Jr. and Hines, A.H. 1992. Interactive effects of salinity and adult extract upon settlement of the estuarine barnacle *Balanus improvisus* (Darwin, 1854). *Journal of Experimental Marine Biology and Ecology* 156: 239-252.
- Doochin, H.D. 1951. The morphology of *Balanus improvisus* Darwin and *Balanus amphitrite niveus* Darwin during initial attachment and metamorphosis. *Bulletin of Marine Science* 1(1): 15-39.
- Dürr, S. and Wahl, M. 2004. Isolated and combined impacts of blue mussels (*Mytilus edulis*) and barnacles (*Balanus improvisus*) on structure and diversity of a fouling community. *Journal of Experimental Marine Biology and Ecology* 306(2): 181-195.
- Foster, B.A. 1970. Responses and acclimation to salinity in the adults of some balanomorph barnacles. *Philosophical Transactions of the Royal Society of London B* 256: 377-400.
- Foster, B.A. and Willan, R.C. 1979. Foreign barnacles transported to New Zealand on an oil platform. *New Zealand Journal of Marine and Freshwater Research* 13(1): 143-149.
- Främmande Arter, 2006. Bay barnacle (*Balanus improvisus*). Fact-sheet available at [http://frammandearter.se/0/2english/pdf/Balanus\\_improvisus.pdf](http://frammandearter.se/0/2english/pdf/Balanus_improvisus.pdf) (accessed on 25 July 2007).
- Furman, E.R. 1990. Geographical variation of *Balanus improvisus* in biochemical and morphometric characters. *Journal*

of the Marine Biological Association of the United Kingdom 70: 721-740.

Furman, E.R. and Yule, A.B. 1990. Self-fertilisation in *Balanus improvisus* Darwin. *Journal of Experimental Marine Biology and Ecology* 144: 235-239.

Furman, E.R., Yule, A.B. and Crisp, D.J. 1989. Gene flow between populations of *Balanus improvisus* Darwin (Cirripedia) in British estuaries. *Topics in Marine Biology. Scientia Marina* 53(2-3): 465-472.

Fyhn, H.J. 1976. Holeuryhalinity and its mechanisms in a cirriped crustacean, *Balanus improvisus*. *Comparative Biochemistry and Physiology* 53A: 19-30.

Gamfeldt, L., Wallén, J., Jonsson, P.R., Berntsson, K.M. and Havenhand, J.N. 2005. Increasing intraspecific diversity enhances settling success in a marine invertebrate. *Ecology* 86(12): 3219-3224.

Gislén, T. 1950. Till kännedomen om invandringen och utbredningen av *Balanus improvisus* vid svenska kuster. *Fauna och Flora* 45: 32-37.

Gollasch, S. and Nehring, S. 2006. National checklist for aquatic alien species in Germany. *Aquatic Invasions* 1(4): 245-269.

Gomoiu, M.-T., Alexandrov, B., Shadrin, N. and Zaitsev, Y. 2002. The Black Sea – a recipient, donor and transit area for alien species. In: *Invasive aquatic species of Europe. Distribution, impacts and management* (eds. E. Leppäkoski, S. Gollasch and S. Olenin), pp. 341-350. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Gouletquer, P., Bachelet, G., Sauriau, P.G. and Noel, P. 2002. Open Atlantic coast of Europe – a century of introduced species into French waters. In: *Invasive aquatic species of Europe. Distribution, impacts and management* (eds. E. Leppäkoski, S. Gollasch and S. Olenin), pp. 276-290. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Grigorovich, I.A., Therriault, T.W. and MacIsaac, H.J. 2003. History of aquatic invertebrate invasions in the Caspian Sea. *Biological Invasions* 5: 103-115.

Hopkins, C.C.E. 2001. Actual and potential effects of introduced marine organisms in Norwegian waters, including Svalbard. Research report 2001-1. Directorate for Nature Management, 53pp.

Iwasaki, K. 2006. Human-mediated introduction of marine organisms in Japan: a review. In: *Assessment and Control of Biological Invasion Risks* (eds. F. Koike, M.N. Clout, M. Kawamichi, M. De Poorter and K. Iwatsuka), pp. 104-112. Shoukado Book Sellers, Kyoto, Japan and IUCN, Gland, Switzerland.

Jensen, K.R. and Knudsen, J. 2005. A summary of alien marine benthic invertebrates in Danish waters. *Oceanological and Hydrobiological Studies* 34, Supplement 1: 137-162.

Kerckhof, F. 2002. Barnacles (Cirripedia, Balanomorpha) in Belgian waters, an overview of the species and recent evolutions, with emphasis on exotic species. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie* 72, Supplement: 93-104.

Kerckhof, F. and Cattrijsse, A. 2001. Exotic Cirripedia (Balanomorpha) from buoys off the Belgian coast. *Senckenbergiana Maritima* 31(2): 245-254.

Kühl, H. 1963. Über die Verbreitung der Balaniden durch Schiffe. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 8: 142-150.

Krüger, P. 1927. Cirripedia Thoracica der Dänischen Gewässer. *Videnskabeligt Meddelelser fra Dansk Naturhistorisk Forening* 84: 11-16.

Laihonen, P. and Furman, E.R. 1986. The site of settlement indicates commensalism between bluemussel and its epibiont. *Oecologia* 71: 38-40.

Lang, W.H., Forward, R.B., Jr. and Miller, D.C. 1979. Behavioral responses of *Balanus improvisus* nauplii to light



intensity and spectrum. *Biological Bulletin* 157: 166-181.

Leppäkoski, E. 1999. *Balanus improvisus* (Darwin 1854), Balanidae, Cirripedia. In: *Exotics Across the Ocean – Case Histories on Introduced Species* (eds. S. Gollasch, D. Minchin, H. Rosenthal and M. Voigt), pp. 49-54. Logos Verlag, Berlin.

Leppäkoski, E. and Olenin, S. 2000. Non-native species and rates of spread: lessons from the brackish Baltic Sea. *Biological Invasions* 2: 151-163.

Minchin, D. 2007. A checklist of alien and cryptogenic aquatic species in Ireland. *Aquatic Invasions* 2(4): 341-366.

Moore, H.B. and Frue, A.C. 1959. The settlement and growth of *Balanus improvisus*, *B. eburneus* and *B. amphitrite* in the Miami area. *Bulletin of Marine Science* 9(4): 421-440.

Nasrolahi, A. 2007. Larval settlement of the barnacle, *Balanus improvisus* Darwin (1854) under different food concentration, substratum, light period, salinity, cyprid density and cyprid age. *Pakistan Journal of Biological Sciences* 10(13): 2231-2236.

O'Connor, N.J. and Richardson, D.L. 1996. Effects of bacterial films on attachment of barnacles (*Balanus improvisus* Darwin) larvae: laboratory and field studies. *Journal of Experimental Marine Biology and Ecology* 206: 69-81.

Olszewska, A. 2000. *Mya arenaria* L., a new and unusual substratum for *Balanus improvisus* Darwin. *Oceanologia* 42(1): 119-121.

Ovsyannikova, L.I. 2008. Barnacles in benthic communities of the inner part of Amursky Bay (Sea of Japan). In: *Ecological studies and the state of the ecosystem of Amursky Bay and the estuarine zone of the Razdolnaya River (Sea of Japan)* (eds. K.A. Lutaenko and M.A. Vaschenko), Vol 1, pp. 207-222. Vladivostok, Dalnauka.

Pitombo, F.B. 2004. Phylogenetic analysis of the Balanidae (Cirripedia, Balanomorpha). *Zoologica Scripta* 33(3): 261-276.

Poulsen, E.M. 1935. De danske farvandes rurer (Balanomorpha og Verrucomorpha). *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening* 99: 5-27.

Rainbow, P.S. 1984. An introduction to the biology of British littoral barnacles. *Field Studies* 6: 1-51.

Rainbow, P.S., Fialkowski, W., Sokolowski, A., Smith, B.D. and Wolowicz, M. 2004. Geographical and seasonal variation of trace metal bioavailabilities in the Gulf of Gdansk, Baltic Sea using mussels (*Mytilus trossulus*) and barnacles (*Balanus improvisus*) as biomonitors. *Marine Biology* 144: 271-286.

Rainbow, P.S., Smith, B.D. and Lau, S.S.S. 2002. Biomonitoring of trace metal availabilities in the Thames estuary using a suite of littoral biomonitors. *Journal of the Marine Biological Association of the United Kingdom* 82: 793-799.

Rainbow, P.S., Wolowicz, M., Fialkowski, W., Smith, B.D. and Sokolowski, A. 2000. Biomonitoring of trace metals in the Gulf of Gdansk, using mussels (*Mytilus trossulus*) and barnacles (*Balanus improvisus*). *Water Research* 34(6): 1823-1829.

Riedel, F., Audzijonytė, A. and Mugue, N. 2006. Aliens associating with Caspian Sea endemic bivalves. *Biological Invasions* 8: 1067-1071.

Ross, A. and Jackson, C.G., Jr. 1972. Barnacle fouling of the ornate diamondback terrapin, *Malaclemys terrapin macropilota*. *Crustaceana* 22(2): 203-205.

Ross, P.M., Burrows, M.T., Hawkins, S.J., Southward, A.J. and Ryan, K.P. 2003. A key for the identification of the nauplii of common barnacles of the British Isles, with emphasis on *Chthamalus*. *Journal of Crustacean Biology* 23(2): 328-340.

Sjögren, M., Dahlström, M., Göransson, U., Jonsson, P.R. and Bohlin, L. 2004. Recruitment in the field of *Balanus*

improvisus and *Mytilus edulis* in response to the antifouling cyclopeptides baretin and 8,9-dihydrobaretin from the marine sponge *Geodia barretti*. *Biofouling* 20(6): 291-297.

Sjögren, M., Dahlström, M., Hedner, E., Jonsson, P.R., Vik, A., Gundersen, L.-L. and Bohlin, L. 2008. Antifouling activity of the sponge metabolite agelasine D. and synthesised analogs on *Balanus improvisus*. *Biofouling* 24(4): 251-258.

Sneli, J.-A. 1972. Distribution of *Balanus improvisus* along the Norwegian coast. *Rhizocrinus* 1(7): 1-6.

Southward, A.J. 1957. On the behaviour of barnacles III. Further observations on the influence of temperature and age on cirral activity. *Journal of the Marine Biological Association of the United Kingdom* 36: 323-334.

Stephensen, K. 1933. Havedderkopper (Pycnogonida) og rankefødder (Cirripedia). *Danmarks Fauna* 38: 1-158.

Streftaris, N., Zenetos, A. and Papathanassiou, E. 2005. Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. *Oceanography and Marine Biology: An annual Review* 43: 419-453.

Toth, G.B. and Lindeborg, M. 2008. Water-soluble compounds from the breadcrumb sponge *Halichondria panicea* deter attachment of the barnacle *Balanus improvisus*. *Marine Ecology Progress Series* 354: 125-132.

Wahl, M., Hay, M.E. and Enderlein, P. 1997. Effects of epibiosis on consumer-prey interactions. *Hydrobiologia* 355: 49-59.

Wonham, M.J. and Carlton, J.T. 2005. Trends in marine biological invasions at local and regional scales: the Northeast Pacific Ocean as a model system. *Biological Invasions* 7: 369-392.

Zega, G., Pennati, R., Dahlström, M., Berntsson, K., Sotgia, C. and de Bernardi, F. 2007. Settlement of the barnacle *Balanus improvisus*: The roles of dopamine and serotonin. *Italian Journal of Zoology* 74(4): 351-361.

Zenetos, A., Çinar, M.E., Panucci-Papadopoulou, M.A., Harmelin, J.G., Furnari, G., Andaloro, F., Bellou, N., Streftaris, N. and Zibrowius, H. 2005. Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science* 6(2): 63-118.

Zvyagintsev, A.Y. 2003. Introduction of species into the northwestern Sea of Japan and the problem of marine fouling. *Russian Journal of Marine Biology* 29 (Supplement 1): S10-S21.