

NOBANIS - Invasive Alien Species Fact Sheet

Anguillicoloides crassus

Authors of this species fact sheet: Viktoras Didžiulis, Catalogue of Life, School of Biological Sciences, Harborne Building, University of Reading, UK, RG6 6AS, viktorasdi@gmail.com

Bibliographical reference – how to cite this fact sheet:

Didžiulis, V. (2013): NOBANIS – Invasive Alien Species Fact Sheet – *Anguillicoloides crassus*. – From: Online Database of the European Network on Invasive Alien Species – NOBANIS www.nobanis.org, Date of access x/x/201x.

Species description

Scientific names: *Anguillicoloides crassus*, Kuwahara, Niimi and Itagaki 1974

Synonyms: -

Common names: Eel swimbladder nematode (GB), Aal-Schwimmblassenwurm (AT), krevnatka úhoří (CZ), ålens svømmeblærenematode (DK), Simblåsemask eller Ålnematod (SE), Schwimmblassenwurm (DE), Angwilikola (PL), Sundmagaormur (IS).



Fig 1. *Anguillicoloides crassus*, photo by Marianne Køie.

Species identification

Adult *Anguillicoloides crassus* nematodes have a body covered by a soft, wrinkled outer cuticle. The circular mouth-opening is small surrounded by 4 dorsolateral and ventrolateral cephalic papillae and 2 small lateral amphids. The anterior rim of the buccal capsule bears one row of 22, 24, 26 or 28 circumoral teeth. The oesophagus appears strongly muscular, consisting of 3 lobes, expanding at its posterior half; anteriorly, the oesophagus forms 6 slightly elevated, rounded lobes protruding into the buccal cavity. The valvular apparatus of the oesophagus is well developed. Male worms have 6 pairs of caudal papillae; 2 praeanal, 1 or 2 adanal and 2 or 3 postanal. The overall body length of males is 20-60 mm with females measuring 47-71.5 mm. Body width of males is .9-2.8 mm and 3-5.6 mm for females (Taraschewski *et al.* 1987). There is a prominent cone-shaped vulva in the posterior part of the body. Mature females with white uterus occupying most of the body, containing numerous eggs, developing embryos and 250 µm long sheathed larvae (Moravec 1994).

Native range

Anguillicoloides crassus originate in Southeast Asia, and its natural host is the Asian eel, *Anguilla japonica* (Kuwahara *et al.* 1974; Taraschewski *et al.* 1987). Related species are also found in New Zealand and Australia (van Banning 1991, Moravec and Taraschewski 1988, Køie 1991).

Alien distribution

History of introduction and geographical spread

Anguillicoloides crassus probably arrived in Europe with eels imported from south-eastern Asia (Höglund *et al.* 1989), possibly from Taiwan at about 1980 (Køie 1991). Within a few years, heavy infestations of the European eel (*Anguilla anguilla*) were reported. It was found in open water of the Weser-Ems River region in Northern Germany and in North Italian culture facilities in 1982 (Neumann 1985, Peters and Hartmann 1986). This nematode successfully colonised most European countries throughout the 1980s, especially in freshwater environments (Ashworth and Blanc 1997), see table 2. The species was found in Germany in 1982 (Køie 1991). In Denmark, the parasite was first observed in 1986 (Køie 1988), in Sweden in 1987 (Höglund *et al.* 1989). The first records for Austria date back to 1987 (Mondsee) and 1988 (Neusiedler See) (Jagsch & Rydlo 1993, Konecny & Wais 1994), where it is still present (Schabuss *et al.* 2005). In Norway, the first infected eels were caught in 1993 (Mo and Steien 1993). In Latvia *A. crassus* was first discovered in lake Usmas and river Venta in 1998, and the parasite was found in all of the examined eels (Kirjusina and Vismanis 2004). *A. crassus* was imported to Estonia with young eels in the middle of the 1980s. In the early 1990s *A. crassus* was distributed practically all over the coastal zone of Estonia (Henn Ojaveer, pers. comm.).

The first observation of *A. crassus* in Sweden was in the artificially heated waters outside the nuclear power-plant Oskarshamn, on the east coast. The parasite rapidly became well established not only in thermal discharge areas but also in other brackish waters, as well as in some freshwater lakes (Wickström *et al.* 1998). In comparison with localities affected by thermal discharges, the infection rate in monitored inland (two sites), and coastal (one site) eel stocks has been much slower. As in Denmark, dispersal by man is believed to have greatly contributed to the further dispersal of the species. According to Wickström *et al.* (1998), there are strong indications that the parasite was introduced into several freshwater lakes as a result of stocking infected wild eels caught on the Swedish west coast.

In Denmark, *A. crassus* was probably first introduced to the brackish Ringkøbing Fjord through trade with the Netherlands. Its further dispersal was probably mediated by man, rather than by natural migration (Boëtius 1989). According to Boëtius (1989) almost all of the [early] infections

can be traced back to the Ringkøbing Fjord (Jutland). Introduction to Denmark probably took place sometime before the first observation in 1986, and within a few years, the parasite was present in several other locations in Jutland and on Zealand.

A. crassus was probably introduced to Estonia with young eels (SI 20-30 cm) from Germany in autumn 1989. In Lake Võrtsjärv the mean infestation rate was 44% (range in 22-71% in different length groups, $n= 141$, SI 50-91 cm) in 1994-97. The mean number of *A. crassus* per infested eel was 24, the maximum 92. (Ojaveer *et al.* 2003)

First record	Country	Reference
1982	Germany	Neumann, 1985
1985	Belgium	Belpaire et al., 1989a
1985	France	Dupont & Petter, 1988
1985	Holland	Van Banning et al., 1985
1986	Denmark	Koie, 1988
1987	Spain	Belpaire et al., 1989a
1987	England	Kennedy & Fitch, 1990
1987	Italy	Canestri-Trotti, 1987
1988	Sardinia (Italy)	Moravec, 1992
1988	Poland	Koops & Hartmann, 1989
1988	Estonia	Kangur, 1994
1988	Sweden	Hellstrom et al., 1988
1988	Greece	Belpaire et al., 1989a
1988?	Egypt	Koops & Hartmann, 1989
1988	Austria	Konecny & Wais, 1993
1990	Hungary	Székely et al., 1991
1991	Czech Republic	Moravec, 1992
1992	Belarus	After Moravec, 1992
1992	Yugoslavia	After Höglund & Thomas, 1992
1992	Portugal	Cruz et al., 1992
1993	Russia, Kaliningrad Oblast	Zaostrovceva, 1993
1993	Norway	Mo & Steien, 1994
1994	Latvia	Vismanis et al., 1999
1994	Tunisia	Maamouri et al., 1999
1994	Marocco	El Hilali et al., 1996
1995	Macedonia	Cakic et al., 2002
1998	Lithuania	Bacevicus, 2004
1998	Ireland	Evans & Matthews, 1999
1999	Algeria	After Nabil et al., 2009
2002	Corsica (France)	Ternengo et al., 2005
2002	Turkey	Genc et al., 2005
2004	Scotland	Lyndon & Pieters, 2005
2007	Finland	Jakob et al., 2009b

Table 1. First records of *Anguillicoloides crassus* in European countries. From Weclawski, U., 2012

Pathways of introduction

The main dispersal vector is believed to be the uncontrolled intercontinental transfer of live eels, including stocking in natural waters (Køie 1991).

Alien status in region

Since its introduction in 1982, *Anguillicoloides crassus* has become firmly established in eel populations in fresh and brackish waters in most European countries (Køie 1991, Eno *et al.* 1997), see also table 2.

In Germany, eel populations are infected already from 65% to 90% with *A. crassus* (Deutscher Bundestag 2004). In eels of the river Rhine the average infection intensities amounted to 6 nematodes per swim-bladder (Wurtz *et al.* 1998). Surveys in other aquatic habitats even revealed average infection intensities of up to 12 nematodes per swimbladder (Leuner 2004).

Country	Not found	Not established	Rare	Local	Common	Very common	Not known
Austria				X			
Czech Republic				X			
Denmark					X		
Estonia					X		
European part of Russia				X			
Finland				X			
Faroe Islands	X						
Germany						X	
Greenland	X						
Iceland	X						
Ireland				X			
Latvia				X			
Lithuania				X			
Netherlands				X			
Norway			X				
Poland					X		
Sweden						X	

Table 2. The frequency and establishment of *Anguillicoloides crassus*, please refer also to the information provided for this species at www.nobanis.org/search.asp. Legend for this table: **Not found** – The species is not found in the country; **Not established** - The species has not formed self-reproducing populations (but is found as a casual or incidental species); **Rare** - Few sites where it is found in the country; **Local** - Locally abundant, many individuals in some areas of the country; **Common** - Many sites in the country; **Very common** - Many sites and many individuals; **Not known** – No information was available.

Ecology

Habitat description

Despite the fact that this is an introduced species, *A. crassus* has been able to find a range of suitable host organisms in its new environment. In addition to various species of marine and freshwater fishes, aquatic snails, amphibians and insects can serve as hosts for *A. crassus* (Moravec and Skorikova 1998). Intermediate hosts include copepods and ostracods (Bauer 1998). The European eel also seems to be more susceptible to the parasite than are their original hosts (Køie

1991). Once introduced into a lake or river it may spread rapidly among the eel population. *Anguillicoloides crassus* is a successful colonizer due to its large production of eggs and low specificity regarding intermediate hosts. Levels of infestation have been recorded to rise from 10% to 50% within a year (Belpaire *et al.* 1989, Koops and Hartmann 1989)

Laboratory investigations have shown that adult parasites and eggs can remain viable in seawater, although hatching off the egg is reduced. The survival of free-living larvae is maximal in freshwater and decline with increasing salinity (Kennedy and Fitch 1990). According to Svedäng (1996) the low number of infected eels on the Swedish west coast compared to the Baltic Sea supports the view that the spread of *A. crassus* is prevented by salinities above 8-10 psu. Infected paratenic hosts in brackish or marine environments, such as ruffe (*Gymnocephalus cernuus*), smelt (*Osmerus eperlanus*), and black goby (*Gobius niger*) can, however, serve as dispersal vectors in marine environments, if they are consumed by eels. Other studies have shown that larval development is significantly retarded at low temperatures and that adult worms show increased mortality and decreased growth and reproduction at low temperatures (Knopf *et al.* 1998). According to Knopf *et al.* (1998) the latter supports the hypothesis that the spread of *A. crassus* in boreal regions is restricted by temperature regimes. On the other hand, increasing temperature will have a negative effect on the establishment of the parasites on intermediate hosts (copepods) (Ashworth *et al.* 1996).

Reproduction and life cycle

This parasite possess several attributes of a successful coloniser such as high dispersal ability, long survival of the free larval stage, high fecundity (large production of eggs), high reproductive rate, and great receptivity of host organisms (Ashworth and Blanc 1997).

A. crassus is ovoviviparous, *i.e.* the eggs are hatched inside the female. The first larval stage emerges within the swimbladder of the eel, where the larva matures to its second stage. By way of the pharynx, the larva reaches the eels' intestine through which it is excreted into the water. The second stage can survive for considerable time in the water, waiting to be consumed by a suitable paratenic or intermediate host, *e.g.* a copepod. Various aquatic non-crustacean invertebrates *i.e.* insects, snails, and vertebrates, amphibians and fish may act as the intermediate host, which is not needed for the development of the parasite, but nonetheless serve to maintain its lifecycle (paratenic hosts) (Kennedy and Finch 1990, Thomas and Ollevier 1992, Moravec and Skorikova 1998). The larvae migrate through the intestinal wall into the swimbladder. There they moult to fourth stage, then molt once again and develop into adults. The life-cycle can be completed in two months in the laboratory (De Charleroy *et al.* 1990).

There is no obvious seasonal cycle in either prevalence or mean intensity of *A. crassus* in the European eel (Thomas and Ollevier 1992, Kelly *et al.* 2000).

Dispersal and spread

In Denmark, as in most of Europe, this eel parasite occurs mainly in freshwater, but it is also found in the brackish Ringkøbing Fjord, which is also believed to be the first point of introduction (Køie 1991). In freshwater it is present *i.a.* in Jutland and on Zealand (Denmark). It is also present in the Poland -Vistula Lagoon and the Pomeranian lakes (Grawinski 1994, Rolbiecki *et al.* 1996). In Sweden, the parasite is firmly established at several locations on the Swedish east coast (Svedäng 1996). It was first discovered in high numbers at sites affected by cooling water discharges from a nuclear power plant (Höglund and Thulin 1994, Höglund and Andersson 1993), but it is no longer limited to areas influenced by thermal discharges. Infected eels are also found along the Swedish west coast, but the levels of infestation are much lower (Svedäng 1996). In Norway, infected eels have been caught just north of the Norwegian/Swedish border (Mo and Steien 1993). *A. crassus* is

not present in Finnish inland waters according to Køie (1991). No further information on its presence (or absence) in Finland since the publication of that paper (1991) has been found.

Due to geothermal activity on Iceland, the all year temperature of some discrete small rivers/ponds is sufficiently warm to support the establishment of *A. crassus*. Iceland's natural isolation acts as a barrier for a natural introduction of alien parasites (the prevention is not absolute as witnessed by the present freshwater fauna) (Kristmundsson and Helgason 2007).

In cultured populations with high densities of eels 100% of the eels may be infected. This can also be the case in wild populations in inland waters (van Banning 1991). The level of infestation in marine and brackish waters seems to be lower, but levels of more than 60% have been reported from brackish waters in Sweden (Höglund *et al.* 1989).

The phenomenon that the prevalence and intensity of infection is high shortly after the introduction of a parasite in a water system followed by stabilization at a lower level has been observed for *A. crassus*. In Lake Arresø in Denmark the prevalence was 3-7% in 1987-1988 (Køie 1988, Boëtius 1989). In 1992 it was close to 100%, but in 2003 the prevalence was less than 50% (Køie, unpubl. obs.). The mean intensity of infection showed a similar development. A similar levelling off in the infection spread of *A. crassus* was found in a canal in southern France. Repeated infections might have made the swimbladder unsuitable for further *A. crassus* establishment (Lefebvre and Crivelli 2004)

Impact

Affected habitats and indigenous organisms

Adverse effects on wild and cultured eels may occur when the level of infestation is high. Damage caused through inflammation of the swimbladder and secondary bacterial infections may decrease growth and increase mortality in the eel (van Banning 1991).

Damage is caused both by larvae, which live in the tissue of the swimbladder, and adults and pre-adults, which suck the blood of the host. Pathological reactions of the tissues include inflammation, necrosis, and scarring. Of all eels examined in a two year German study, 28% showed pathological alterations of the swimbladder (Wurtz *et al.* 1998). Repeated infestations may eventually reduce the swimbladder to a “non-functional mass of tissues” (Hartmann and Nellen 1997). Concern has been raised that massive damage to the swimbladder may make the eels unable to migrate back to their breeding grounds in the Sargasso Sea. This could have a severe impact on the recruitment of eels and thus on the future of the European eel population (Hartmann and Nellen 1997).

Kelly *et al.* (2000) found that there were no major differences in hormonal, metabolic or osmoregulatory status of eels infected with up to 15 blood-feeding adults in the swimbladder lumen and up to 25 larvae in the swimbladder wall compared with wild uninfected eels. Gollock *et al.* (2005) found no significant changes in mean erythrocyte haemoglobin concentration in uninfected and infected eels. They suggested that acute temperature increase alone is unlikely to cause significant mortality of eels infected with *A. crassus*. However, the effects of chronic increases in temperature in combination with other factors such as toxicants and hypoxia are unknown. In lake Balaton, Hungary, mass mortality of eels infected with *A. crassus* was concluded to have resulted from high infection rates combined with high temperatures (Molnár *et al.* 1991).

The nematode can have negative effects on the condition of eels, especially those under stress. In cultured eels loss of appetite and vitality resulting in emaciation has been reported (van Banning 1991). Effects on wild populations are more difficult to study. Laboratory studies of stress factors that appear in the wild, such oxygen stress, may have a bearing on natural population as well. Experiments with decreasing oxygen levels have demonstrated that the eels most severely affected by *anguillicoloides* are the first to die, and the author concludes that infection by *anguillicoloides* impairs the eels' natural resistance and, under unfavourable environmental conditions, may lead to their death (Molnár 1993). Loss of vitality may also lead to greater susceptibility to infections, and to an increased risk of being preyed upon.

Genetic effects

Not known.

Human health effects

Not known.

Economic and societal effects (positive/negative)

Infected eels have reduced swimming ability (Sprengel and Luchtenberg 1991). It is thus likely that heavily infected eels have an increased risk of being preyed upon by *e.g.* cormorants.

The parasite has caused mortalities in Danish eel farms (Køie 1991). Even if the eels do not die, infection may cause a reduction of the growth rate, and thus in the economic output of the eel farms. It is unknown whether damage to the swimbladder may prevent the spawning migration to the Sargasso Sea. Uninfected eels are found in the marine environment, so *A. crassus* has probably only a limited effect, if any, on the European eel fisheries. Overfishing and pollution are greater risks.

Management approaches

Prevention methods

For cultured fish drug treatment, either by bathing the fish or administered with the feed, has been tested, with varying degrees of success (Taraschewski *et al.* 1988).

Eradication, control and monitoring efforts

Since dispersal by man has been identified as the main vector for the continued spread of *A. crassus* in Europe, monitoring and strict quarantine measures will be necessary to control its further spread. No dedicated programmes are running so far.

Information and awareness

This species presents a serious threat, therefore it is always in focus in the most scientific and popular publications relating to the general topic of invasive species and their spread in the Europe. However, interested parties are mostly scientists and aquaculturists. Any information on further spread and development of prevalence and intensities of infection is of interest as well as observations on diseased eels in relation to their natural environment.

Recommendations or comments from experts and local communities

Since spread between localities is generally through transport of infected eels, uncontrolled movements of infected eels should be avoided.

References and other resources

Contact persons

Robert Konecny (AT), Environment Agency Austria, Spittelauer Lände 5, A-1090 Wien, Austria;
Phone: +43 1 31304 3581; E-mail: robert.konecny@umweltbundesamt.at

Stefan Nehring (DE), AeT umweltplanung, Bismarckstrasse 19, D-56068 Koblenz, Germany;
Phone: +49 261 1330398; E-mail: nehring@aet-umweltplanung.de

Marianne Køie (DK) Marine Biological Laboratory, University of Copenhagen, Strandpromenaden 5, DK-3000 Helsingør, Denmark, Phone: +45 35 32 19 81, E-mail: mkoie@bi.ku.dk,

Henn Ojaveer (EE), Estonian Marine Institute, University of Tartu, Vana-Sauga 28, 80031 Parnu, Estonia, Tel + 372 44 33800; + 372 51 58328, Fax + 372 44 33800, E-mail: henn.ojaveer@ut.ee

Árni Kristmundsson (IS) Fish disease laboratory, Institute for Experimental Pathology, University of Iceland, Keldur v/Vesturlandsveg, IS-112 Reykjavik, Iceland, Tel. +354 5855100, Fax +354 5673979, E-mail: arni@hi.is

Agnis Ozoliņš (LV) Latvian Fish Resources agency, Laboratory of Aquaculture, Daudavgrīvas 8, LV 1048, Riga, phone: +371 7610088, e-mail: agnis.ozolins@latzra.lv

Leszek Rolbiecki (PL) University of Gdańsk, Faculty of Biology, Geography and Oceanology, Chair of Zoology of Invertebrates, Al. Marszałka Piłsudskiego 46; PL-81-378 Gdynia, Poland, Phone: +48 58 6601600, fax +48 58 6601676; E-mail: rolbieck@sat.ocean.univ.gda.pl

Håkan Wickström (SE) Swedish Board of Fisheries, Institute of Freshwater research, Stångholmsvägen2, SE-178 93 Drottningholm, Sweden tel.: +46 (0) 86990607, E-mail: hakan.wickstrom@fiskeriverket.se

Links

EELREP project [papers and manuscripts](#)

Aquatic alien species in German inland and coastal waters ([database](#))

References

- Ashworth, S. T., Kennedy, C. R., and Blanc, G., 1996. Density-dependent effects of *Anguillicola crassus* (Nematoda) within and on its copepod intermediate hosts. *Parasitology* 113(3), 303-309.
- Ashworth, S. T., and Blanc, G., 1997. *Anguillicola crassus*, a recently introduced aggressive colonizer of European eel stocks. - *Bulletin Francais de la Peche et de la Pisciculture* 344/345: 335-342.
- Bacevicus, E. (2004). The new data on ecology of alien fish species and fish metazoan parasites in the Curonian Lagoon and Lithuanian coastal zone. *Baltic – the Sea of Aliens*. Gdynia, Poland. Book of Abstracts. 17–18.
- Bauer, O. N., 1998. A new parasitic nematode from the genus *Anguillicola* (Dracunculioidea: Anguillicolidae) in Palaearctic fish. - *Parazitologiya* 32: 59-65.
- Belpaire, C., De Charleroy, D., Thomas, K., van Damme, P. and Ollevier, F. 1989. Effects of eel restocking on the distribution of the nematode *Anguillicola crassus* in Flanders, Belgium. - *J. Appl. Ecol.* 5: 151-153.
- Boëtius, I., 1989. Preliminary report on the occurrence of *Anguillicola* in some Danish fresh- and seawater areas. - *Internationale Revue der gesamte Hydrobiologie* 75: 889.

- Cakić, D. P., Stojanovski, S., Kulišić, Z., Hristovski, N. & Lenhardt, M. (2002). Pojava *Anguillicola crassus*, nematoda: Dracunculoidea, kod jegulje iz Ohridskog jezera, Makedonija. *Acta veterinaria* 52, 163–168.
- Canestri-Trotti, G. (1987). Occurrence of the Nematode *Anguillicola crassa* (Kuwahara, Niimi & Itagaki, 1974) in eels from the Po delta, Italy. *Bulletin-European Association of Fish Pathologists* 7(5), 109–111.
- Cruz, P., Silva, E., Grazina Freitas, M. S. & Carvalha-Varela, M. (1992). First report of *Anguillicola crassus* in the European eel in Portugal. *Bulletin-European Association of Fish Pathologists* 12, 154–154.
- De Charleroy D, L. Grisez, K. Thomas, C. Belpaire, and F. Ollivier. 1990. The life cycle of *Anguillicola crassus*. – *Dis. Aquat. Org.* 8: 77-84.
- Deutscher Bundestag 2004. Zukunft von Aalzucht und Aalfischerei in Deutschland. - Antwort der Bundesregierung auf eine Kleine Anfrage, Deutscher Bundestag Drucksache 15/2929: 1-8.
- Dupont, F. & Petter, A. J. (1988). *Anguillicola*, a multispecific epizootic in Europe: appearance of *Anguillicola crassa* (Nematoda, Anguillicolidae) in the European eel *Anguilla anguilla* in Camargue, southern France. *Bull. Fr. Pêche Piscic* 308, 38–41.
- El Hilali, M., Yahyaoui, A., Sadak, A., Maachi, M. & Taghy, Z. (1996). Premières données épidémiologiques sur l'anguillicolose au Maroc. *Bulletin Français de la Pêche et de la Pisciculture* 340, 57–60.
- Eno, C. N., Clark, R. A., and Sanderson, W. G., 1997. Non-native marine species in British waters: a review and directory. - Joint Nature Conservation Committee Peterborough.
- Evans, D. W. & Matthews, M. A. (1999). *Anguillicola crassus* (Nematoda, Dracunculoidea); first documented record of this swim bladder parasite of eels in Ireland. *Journal of fish biology* 55(3), 665–668.
- Genc, E., Şahan, A., Altun, T., Cengizler, İ. & Nevsat, E. (2005). Occurrence of the swim bladder parasite *Anguillicola crassus* (Nematoda, Dracunculoidea) in European eels (*Anguilla anguilla*) in Ceyhan River, Turkey. *Turkish Journal of Veterinary and Animal Sciences* 29, 661–663.
- Gollock M.J., C.R. Kennedy, and J.A. Brown. 2005. Physiological responses to acute temperature increase in European eels *Anguilla anguilla* infected with *Anguillicola crassus*. – *Dis. Aquat. Org.* 64: 223-228.
- Grawiński E., 1994. Occurrence of *Anguillicola crassa* nematode in eel (*Anguilla anguilla*) from Vistula Lagoon and Pomeranian lakes, XVII Zjazd Polskiego Towarzystwa Parazytologicznego, 15-16 September, 1994, Gdynia, Poland: 55.
- Hartmann, F., and Nellen, W., 1997. The status of the European eel, *Anguilla anguilla*, after the invasion of the introduced swim bladder parasite, *Anguillicola crassus*, using the eel population in the lower Elbe as an example. - In ICES Annual Science Conference, (Theme Session, 9 pp.). Baltimore, MD (USA), 25 Sep-3 Oct 1997: ICES (International Council for the Exploration of the Sea), Copenhagen, Denmark.
- Hellstrom, A., Ljungberg, O. & Bornstein, S. (1988). *Anguillicola*, en ny ålparasit i Sverige. *Svensk Veterinartidning* 40 (4) 211–213.
- Höglund, J., Andersson, J., and Thulin, J., 1989. Den asiatiska ålparasiten *Anguillicola* nu etablerad i svenska kustvatten - Rapport 3616. Naturvårdsverket, Laboratoriet för miljökontroll.
- Höglund J., and J. Andersson, 1993. Prevalence and abundance of *Anguillicola crassus* in the European eel (*Anguilla anguilla*) at a thermal discharge site on the Swedish coast. – *J. Appl. Ichthyol.* 9: 115-122.
- Höglund, J., and Thulin, J., 1994. Ny sjukdom hos ål – här för att stanna - *Forskning och Framsteg* 2: 25-29.
- Jakob, E., Walter, T. & Hanel, R. (2009b). A checklist of the protozoan and metazoan parasites of European eel (*Anguilla anguilla*): checklist of *Anguilla anguilla* parasites. *Journal of Applied Ichthyology* 25, 1-49.
- Kangur, A. (1994). Uus liik Vortsjaerve kalaparasitide hulgas. *Eesti loodus* 6, 189.
- Kelly, C.E., C.R. Kennedy, and J.A. Brown. 2000. Physiological status of wild European eels (*Anguilla anguilla*) infected with the parasitic nematode, *Anguillicola crassus*. – *Parasitology* 120: 195-202
- Kennedy, C. R., and Fitch, D. J., 1990. Colonization, larval survival and epidemiology of the nematode *Anguillicola crassus*, parasitic in the eel, *Anguilla anguilla*, in Britain. - *Journal of Fish Biology* 36: 117-131.
- Kirjusina M. and Vismanis K., 2004. Freshwater and marine fish parasites in Latvia. Classed catalogue. GosNIORKh, St. Peterburg, Russia: 71.
- Knopf, K., Wurtz, J., Sures, B., and Taraschewski, H., 1998. Impact of low water temperature on the development of *Anguillicola crassus* in the final host *Anguilla anguilla*. - *Diseases of Aquatic Organisms* 33: 143-149.
- Konecny, R., & Wais, A., 1994. Occurrence of *Anguillicola crassus* in eels of Lake Neusiedl, Austria. - *Folia Parasitologia* 40: 327.
- Koops, H., & Hartmann, F., 1989. *Anguillicola* infestations in Germany and in German eel imports. *Journal of Applied Ichthyology*, 1: 41-45.
- Køie, M., 1988. Parasites in European eel *Anguilla anguilla* (L.) from Danish freshwater, brackish and marine localities. - *Ophelia* 29: 93-118.
- Køie, M., 1991. Swimbladder nematodes (*Anguillicola* spp.) and gill monogeneans (*Pseudodactylogyrus* spp.) parasitic on the European eel (*Anguilla anguilla*). - *J. Cons. int. Explor. Mer* 47: 391-398.
- Kristmundsson A. & Helgason, S. 2007.. Parasite communities of eels (*Anguilla anguilla* L.) in

- freshwater and marine habitats in Iceland in comparison with other parasite communities of eels in Europe. *Folia Parasitologica* **54**: 141-153
- Kuwahara, A., A. Niimi, H. Itagaki. 1974. Studies On The Nematode Parasitic On The Air Bladder Of Eel. I. Description Of *Anguillicola Crassa* N. Sp. (Philometridae, Anguillicolidae). *Japanese Journal Of Parasitology* **23**: 275-279.
- Lefebvre F.S., and A.J. Crivelli. 2004. Anguillicolosis: dynamics of the infection over two decades. *Dis. Aquat. Org.* **62**: 227-232.
- Leuner, E. 2004. Untersuchungen an Schwimmblasen von Aalen. Bayerische Landesanstalt für Landwirtschaft, Institut für Fischerei Jahresbericht 2003: 25.
- Lyndon, A. R. & Pieters, N. (2005). First record of the eel swim bladder parasite *Anguillicola crassus* (Nematoda) from Scotland. *Bulletin-European Association of Fish Pathologists* **25**(2), 82.
- Maamouri, F., Gargouri, L., Ould Daddah, M. & Bouix, G. (1999). Occurrence of *Anguillicola crassus* (Nematode, Anguillicolidae) in the Ichkeul Lake (Northern Tunisia). *Bulletin-European Association of Fish Pathologists* **19**, 17-19.
- Mo, T. A., and Steien, S. H., 1993. First observation of the eel swimbladder nematode *Anguillicola crassus* in Norway. *Bulletin of the European Association of Fish Pathologists* **14**: 163-164.
- Molnár, K., C. Székely, and F. Baska. 1991. Mass mortality of eel in Lake Balaton due to *Anguillicola crassus* infection. *Bull. Eur. Assoc. Fish Pathol.* **11**: 211-212.
- Molnár, K., 1993. Effect of decreased oxygen content on eels (*Anguilla anguilla*) infected by *Anguillicola crassus* (Nematoda: Dracunculoidea). *Acta Veterinaria Hungarica* **41**: 349-360.
- Moravec, F., and Skorikova, B., 1998. Amphibians and larvae of aquatic insects as new paratenic hosts of *Anguillicola crassus* (Nematoda: Dracunculoidea), a swimbladder parasite of eels. *Diseases of Aquatic Organisms* **34**: 217-222.
- Moravec, F., & H. Taraschewski. 1988. Revision of the genus *Anguillicola* Yamaguti, 1935 (Nematoda: Anguillicolidae) of the swimbladder of eels, including descriptions of two new species, *A. novaezelandiae* sp. n. and *A. papernai* sp. n. *Folia Parasitologica* **35**: 125-146.
- Moravec F. 1994. Parasitic nematodes of freshwater fishes of Europe, Kluwer Academic publishers, Dordrecht, Boston, London, 273pp.
- Nabil, L., Abderrafik, M. & Boudjema, S. (2009). Biodiversité des Parasites chez *Anguilla anguilla* Linnaeus, 1758 dans le Parc National d'El Kala – Algérie. *European Journal of Scientific Research* **25**(2), 300-309.
- Neumann, W., 1985. Schwimmblasenparasit *Anguillicola* bei Aalen. *Fischer und Teichwirt* **11/1985**: 322.
- Ojaveer, H., Pihu, E., Saat, T. 2003. Fishes of Estonia. Tallinn.
- Peters, G. and F. Hartmann. 1986. *Anguillicola*, a parasitic nematode of the swim bladder spreading among eel populations in Europe. - *Dis. aquat. Org.* **1**: 229-230.
- Rolbiecki L., Grawiński E., Rokicki J., 1996. The occurrence of nematode *Anguillicola crassus* Kuwahara, Niimi et Itagaki 1974 in the swimbladder of eel (*Anguilla anguilla* (L.)) from the Vistula Lagoon, Land-Ocean Interactions in the Coastal Zone, Second International Symposium on: Functioning of Coastal Ecosystems in Various Geographical Regions, 5-7 September, 1996, Sopot, Poland: 55-56.
- Schabuss, M., Kennedy, C.R., Konecny, R., Grillitsch, B., Reckendorfer, W., Schiemer, F. Herzig, A., 2005. Dynamics and predicted decline of *Anguillicola crassus* infection in European eels, *Anguilla anguilla*, in Neusiedler See, Austria. - *Journal of Helminthology* **79**: 159-167.
- Sprengel G., H. Lüchtenberg. 1991. Infection by endoparasites reduces maximum swimming speed of European smelt *Osmerus eperlanus* and European eel *Anguilla anguilla*. *Dis. Aquat. Org.* **11**: 31-35.
- Svedäng, H., 1996. Spridning av simblåsemasken *Anguillicola crassus* hos ål i svenska kustvatten - Kustrapport No. 1996:4. Fiskeriverket, Kustlaboratoriet.
- Székely, C., Lang, M. & Csaba, G. (1991). First occurrence of *Anguillicola crassus* in Hungary. *Bulletin of the European Association of Fish Pathologists* **11**(4), 162-163.
- Taraschewski, H., F. Moravec, T. Lamah, K. Anders, 1987. Distribution And Morphology Of Two Helminths Recently Introduced Into European Eel Populations: *Anguillicola Crassus* (Nematoda, Dracunculoidea) And *Paratenuisentis Ambiguus* (Acanthocephala, Tenuisentidae). *Diseases Of Aquatic Organisms* **3**:167-176.
- Taraschewski H., C. Renner, and H. Mehlhorn. 1988. Treatment of fish parasites. 3. Effects of levamisole HCL, metrifonate, fenbendazole, mebendazole, and ivermectin on *Anguillicola crassus* (nematodes) pathogenic in the air bladder of eels. *Parasitol. Res.* **74**: 281-289.
- Ternengo, S., Levron, C., Desideri, F. & Marchand, B. (2005). Parasite communities in European eels *Anguilla anguilla* (Pisces, Teleostei) from a Corsican coastal pond. *Vie et milieu* **55**(1), 1-6.
- Thomas K., and F. Ollevier, 1992. Paratenic hosts of the swimbladder nematode *Anguillicola crassus*. *Dis. Aquat. Org.* **13**: 165-174.

- van Banning P., 1991. Swimbladder nematode (*Anguillicola crassus*) in the European eel (*Anguilla anguilla*) (ICES Identification Leaflets for Diseases and Parasites of Fish and Shellfish. Leaflet No. 48). International Council for the Exploration of the Sea.
- Vismanis, K., Kirjusina, M. & Rodzins, R. (1999). An eel nematode *Anguillicola crassus* Kuwahara, Nimii et Itagaki, 1974 (Nematoda: Dracunculoidea) in Latvia. *Proceedings of the Fifth International Symposium on Fish Parasites, Czech Republic*.
- Weclawski, U., 2012. Evolutionary divergence of *Anguillicola crassus*, an invasive parasitic swim bladder nematode of eels of the genus *Anguilla*. Dissertation, Fakultät für Chemie und Biowissenschaften, Karlsruher Institut für Technologie (KIT) <http://d-nb.info/1022123831/34>
- Wickström, H., Clevestam, P., and Höglund, J., 1998. The spreading of *Anguillicola crassus* in freshwater lakes in Sweden. *Bulletin Francais de la Peche et de la Pisciculture* 349: 215-221.
- Wurtz, J, Knopf, K, and H. Taraschewski. 1998. Distribution and prevalence of *Anguillicola crassus* (Nematoda) in eels *Anguilla anguilla* of the rivers Rhine and Naab, Germany. *Dis. Aquat. Organ.* 32: 137-143.
- Zaostrovceva, S. (1993). Ekologo-faunisticheskij obzor gelmintofauny ryb Kaliningradskogo Zaliva. Nauchno-tehnicheskaja Konferencija Professorsko-Prepodavatel'skogo Sostava, Aspirantov i Sotrudnikov Kaliningradskogo Tekhnicheskogo Instituta Rybnoj Promyshlennosti i Khozjajstva, Tezisy dokladov 18.

Date of creation/modification of this species fact sheet: 18-12-2006/ 04-06-2013